

Stroke

American Stroke
AssociationSM

JOURNAL OF THE AMERICAN HEART ASSOCIATION

A Division of American
Heart Association



**{alpha}-Galactosidase A Deficiency Leads to Increased Tissue Fibrin Deposition
and Thrombosis in Mice Homozygous for the Factor V Leiden Mutation**

Yuechun Shen, Peter F. Bodary, Fernando B. Vargas, Jonathon W. Homeister, David
Gordon, Kristen A. Ostenso, James A. Shayman and Daniel T. Eitzman

Stroke 2006;37;1106-1108; originally published online Mar 2, 2006;

DOI: 10.1161/01.STR.0000206442.86238.39

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2006 American Heart Association. All rights reserved. Print ISSN: 0039-2499. Online
ISSN: 1524-4628

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://stroke.ahajournals.org/cgi/content/full/37/4/1106>

Subscriptions: Information about subscribing to Stroke is online at
<http://stroke.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters
Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax:
410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

α -Galactosidase A Deficiency Leads to Increased Tissue Fibrin Deposition and Thrombosis in Mice Homozygous for the Factor V Leiden Mutation

Yuechun Shen, Peter F. Bodary, Fernando B. Vargas, Jonathon W. Homeister, David Gordon, Kristen A. Ostenson, James A. Shayman, Daniel T. Eitzman

Background—Factor V Leiden (FVL) is a common genetic risk factor for vascular thrombosis in humans. Fabry disease, an X-linked lysosomal storage disorder attributable to α -galactosidase A (GLA) deficiency, is associated with premature vascular events that may be thrombotic in nature.

Methods and Results—To examine a potential interaction between *FvL* and *Gla* deficiency in vivo, we analyzed tissue fibrin deposition in mice carrying combined mutations in *FvL* and *Gla*. *Gla* deficiency markedly increased tissue fibrin deposition in mice carrying the *FvL* mutation ($0.33 \pm 0.03\%$; $n=7$) compared with *FvL* mutation ($0.14 \pm 0.02\%$; $n=10$; $P<0.0005$).

Conclusions—These observations demonstrate a synergistic interaction between *Gla* deficiency and *FvL* toward tissue fibrin deposition in mice. Concomitant mutations in these genes may increase the penetrance of vascular thrombotic events in humans. (*Stroke*. 2006;37:1106-1108.)

Key Words: fibrin ■ genetics ■ thrombosis

A common point mutation in factor V (factor V Leiden [FVL]; 2% to 7% prevalence for European populations) leads to activated protein C resistance and thrombophilia.¹ Several reports in humans and animals have documented a synergistic risk of thrombosis when FVL is combined with other genetic modifiers of thrombosis.²

Fabry disease is an X-linked disorder that results from deficiency of α -galactosidase A (GLA) enzymatic activity.³ Premature vascular events in Fabry patients suggest a propensity toward vascular thrombosis.⁴ An increased thrombotic response has been shown to occur after arterial injury in *Gla*-deficient mice,⁵ although the effect of *Gla* deficiency in spontaneous thrombosis is unclear.

To determine whether *Gla* deficiency in mice is associated with an increased tendency toward spontaneous thrombosis, compound mutant mice carrying mutations in *Gla* and *FvL* were generated and analyzed for tissue fibrin deposition and thrombosis.

Methods

FvL (*FvL*^{0/0})⁶ mice were crossed to mice deficient in *Gla*⁷ and then genotyped as previously described.⁶⁻⁷ Mice were perfusion fixed with zinc formalin, and sections were stained for fibrin(ogen) as done previously.² Tissue fibrin(ogen) was graded by a blinded observer using Image-Pro Plus software (Media Cybernetics). Each section was quantitated for percentage fibrin staining using automated color detection. Hematoxylin and eosin along with fibrin-stained sections were also reviewed for the presence of thrombus.

Thrombi in each section were counted by an observer blinded to mouse genotype and were defined as organized, fibrin-stained, vascular occlusions.

Values are expressed as mean \pm SEM. The statistical significance of differences between groups was determined by 1-way ANOVA followed by Dunn's post hoc analysis when >2 experimental groups were included. The Student 2-tailed *t* test was performed when only 2 groups were being compared. $P<0.05$ was considered significant.

Results

At 22.3 ± 0.6 months of age, 5 organs from each mouse, including kidney, lung, liver, heart, and brain, were analyzed for fibrin deposition. Mice homozygous for *FvL* (*Gla*⁺⁰ *FvL*^{0/0}) demonstrated increased fibrin deposition (Figure 1A) compared with wild-type mice (*Gla*⁺⁰ *FvL*⁺⁰) as described previously. Fibrin deposition in mice with deficiency of *Gla* (*Gla*⁻⁰ *FvL*⁺⁰) alone was not significantly elevated compared with wild-type (WT) mice (*Gla*⁺⁰ *FvL*⁺⁰). However, in the presence of *FvL*, deficiency of *Gla* (*Gla*⁻⁰ *FvL*^{0/0}) greatly increased tissue fibrin deposition compared with either homozygous *FvL*- or *Gla*-deficient mice (Figure 1A).

To determine the effect of heterozygous *Gla* deficiency, female mice were also analyzed. Homozygous *FvL* mice with deficiency of *Gla* (*Gla*^{-/-} *FvL*^{0/0}) showed significantly increased fibrin staining compared with mice homozygous for *FvL* with WT *Gla* (*Gla*^{+/+} *FvL*^{0/0}). *Gla* heterozygous *FvL* mice (*Gla*^{+/-} *FvL*^{0/0}) were intermediate between *Gla*^{+/+} *FvL*^{0/0} and

Received October 14, 2005; accepted November 9, 2005.

From the Department of Internal Medicine, Division of Cardiovascular Medicine, (Y.S., P.F.B., F.B.V., K.A.O., D.T.E.), Division of Nephrology (J.A.S.), and Department of Pathology (J.W.H., D.G.), University of Michigan Medical Center, Ann Arbor.

Correspondence to Daniel T. Eitzman, MD, 1150 W Medical Center Dr, 7301 MSRB III, Ann Arbor, MI 48109-0644. E-mail deitzman@umich.edu
© 2006 American Heart Association, Inc.

Stroke is available at <http://www.strokeaha.org>

DOI: 10.1161/01.STR.0000206442.86238.39




Figure 1. Analysis of fibrin(ogen) in organs. A, Quantification of fibrin(ogen). In the presence of *FvL* homozygosity, deficiency of *Gla* ($Gla^{-/0} FvL^{Q/Q}$; n=7) greatly increased tissue fibrin deposition compared with homozygous *FvL* ($Gla^{+/0} FvL^{Q/Q}$; n=10) or deficiency of *Gla* mice alone ($Gla^{-/0} FvL^{+/+}$; n=4). B, Heterozygous deficiency of *Gla* mice ($Gla^{+/-} FvL^{Q/Q}$; n=7) were intermediate between homozygous ($Gla^{-/-} FvL^{Q/Q}$; n=7) and WT *Gla* mice ($Gla^{+/+} FvL^{Q/Q}$; n=4). WT mice ($Gla^{+/+} FvL^{+/+}$; n=4) were used as control. * $P < 0.05$; ** $P < 0.0005$. C, Immunohistochemical staining of fibrin(ogen), shown as red. Bar=40 μ m.

$Gla^{-/-} FvL^{Q/Q}$ mice for tissue fibrin deposition (Figure 1B and 1C) but not significantly different from either group.

Thrombi from each of 5 organs were counted from all mice included in the fibrin analysis. No thrombi were observed in WT mice or mice with *Gla* deficiency. Thrombi were only identified in mice homozygous for *FvL* ($Gla^{+/+} FvL^{Q/Q}$ and $Gla^{+/0} FvL^{Q/Q}$; 0.3 ± 0.1 thrombi/mouse) and were present in veins of the kidneys and lungs. However, with concomitant *Gla* deficiency ($Gla^{-/-} FvL^{Q/Q}$ and $Gla^{-/0} FvL^{Q/Q}$), the mean number of thrombi was significantly increased (1.9 ± 0.7 thrombi/mouse; $P < 0.04$). Thrombi were observed in veins of the kidney, lung, and liver and 2 $Gla^{-/0} FvL^{Q/Q}$ mice exhibited organized thrombi in major coronary arteries (Figure 2). During the 22-month observation period, we observed 1 mouse with an apparent spontaneous stroke at 18 months of age that was of the $Gla^{-/-} FvL^{Q/Q}$ genotype. Histologic analysis demonstrated cerebral arterial thrombosis with perivascular inflammation.

Discussion

The *FvL* mouse is a useful model to uncover relevant genetic modifiers of thrombosis. Mice homozygous for the murine *FvL* mutation display activated protein C resistance and spontaneously deposit fibrin in their tissues,⁶ suggesting chronic low-grade thrombin generation. This model has been used previously to unmask the phenotype of antithrombotic genes.²

In the current study, we used the *FvL* mouse model to determine the effect of *Gla* deficiency on spontaneous thrombosis in mice. In the presence of *FvL* homozygosity, deficiency of *Gla* greatly increased fibrin deposition and occlusive thrombus formation compared with mice homozygous for *FvL* or with deficiency of *Gla* alone. This observation suggests that under certain circumstances, *Gla* deficiency leads to increased propensity toward spontaneous thrombosis. Although the mechanism is unclear, a vascular wall defect




Figure 2. Organized occlusive thrombus in a coronary artery. Fibrin(ogen) staining of heart from $Gla^{-/-} FvL^{Q/Q}$ mouse. Thrombus stains red. Bar=60 μ m.

leading to dysregulation of NO with resultant oxidative stress may play a role.⁸⁻⁹

The findings in this study are consistent with a recent human clinical study, which found increased ischemic cerebral lesions in Fabry patients carrying the FVL mutation.¹⁰

Acknowledgments

This work was supported by National Institutes of Health (NIH) grant PO1 HL-57346 to D.E., D.G., and J.H. and NIH grant RO1 DK-055823 to J.S. *FvL* mice were from Drs David Ginsburg and Randal J. Westrick (HHMI, University of Michigan).

References

1. Svensson PJ, Dahlbäck B. Resistance to activated protein C as a basis for venous thrombosis. *N Engl J Med.* 1994;330:517-522.
2. Eitzman DT, Westrick RJ, Bi X, Manning SL, Wilkinson JE, Broze GJ, Ginsburg D. Lethal perinatal thrombosis in mice resulting from the interaction of tissue factor pathway inhibitor deficiency and factor V Leiden. *Circulation.* 2002;105:2139-2142.
3. Brady RO, Gal AE, Bradley RM, Martensson E, Warshaw AL, Laster L. Enzymatic defect in Fabry's disease. Cereamidetrihexosidase deficiency. *N Engl J Med.* 1967;276:1163-1167.
4. Utsumi K, Yamamoto N, Kase R, Takata T, Okumiya T, Saito H, Suzuki T, Uyama E, Sakuraba H. High incidence of thrombosis in Fabry's disease. *Intern Med.* 1997;36:327-329.
5. Eitzman DT, Bodary PF, Shen Y, Khairallah CG, Wild SR, Abe A, Shaffer-Hartman J, Shayman JA. Fabry disease in mice is associated with age-dependent susceptibility to vascular thrombosis. *J Am Soc Nephrol.* 2003;14:298-302.
6. Cui J, Eitzman DT, Westrick RJ, Christie PD, Xu ZJ, Yang AY, Purkayastha AA, Yang TL, Metz AL, Gallagher KP, Tyson JA, Rosenberg RD, Ginsburg D. Spontaneous thrombosis in mice carrying the factor V Leiden mutation. *Blood.* 2000;96:4222-4226.
7. Ohshima T, Murray GJ, Swaim WD, Longenecker G, Quirk JM, Cardarelli CO, Sugimoto Y, Pastan I, Gottesman MM, Brady RO, Kulkarni AB. α -Galactosidase A deficient mice: a model of Fabry disease. *Proc Natl Acad Sci.* 1997;94:2540-2544.
8. Moore DF, Scott LT, Gladwin MT, Altarescu G, Kaneski C, Suzuki K, Pease, Fye M, Ferri R, Brady RO, Herscovitch P, Schiffmann R. Regional cerebral hyperperfusion and nitric oxide pathway dysregulation in Fabry disease: reversal by enzyme replacement therapy. *Circulation.* 2001;104:1506-1512.
9. Bodary PF, Shen Y, Vargas FB, Bi X, Ostenson KA, Gu S, Shayman JA, Eitzman DT. Alpha-galactosidase A deficiency accelerates atherosclerosis in mice with apolipoprotein E deficiency. *Circulation.* 2005;111:629-632.
10. Altarescu G, Moore DF, Schiffmann R. Effect of genetic modifiers on cerebral lesions in Fabry disease. *Neurology.* 2005;64:2148-2150.