

- sine-induced dilatation of pial arteries of cats. *Blood Vessels* 14: 285-293, 1977
17. Lux HD: Extracellular potassium in the CNS: relation to excitability changes. *In* Ingvar D, Lassen N (eds): *Brain Work*. Copenhagen, Munkgaard, pp 172-181, 1975
 18. Lewis DV, Schuette WH: NADH fluorescence and K^+ changes during hippocampal electrical stimulation. *J Neurophysiol* 38: 405-417, 1975
 19. Raberger G, Weissel M, Kraupp O: The dependence of the effects of i. cor. administered adenosine and of coronary conductance on the arterial pH, pCO_2 and buffer capacity in dogs. *Arch Pharmacol* 271: 301-310, 1971
 20. Burnstock G: Purinergic transmission. *In* Iverson LL, Iverson SD, Snyder SH (eds): *Handbook of Psychopharmacology*, vol 5. New York, Plenum Publishing Corporation, pp 131-194, 1975
 21. Holmsen H, Day HJ, Stormorken H: The blood platelet release action. *Scand J Haemat (Suppl 8)*: 1-26, 1969
 22. Kalendovsky Z, Austin JH: "Complicated migraine," its association with increased platelet aggregability and abnormal plasma coagulation factors. *Headache* 15: 18-35, 1975
 23. Norwood ChW, Poole GJ, Moody D: Treatment of experimental delayed cerebral arterial spasm with a beta₂-adrenergic stimulator and a phosphodiesterase inhibitor. *J Neurosurg* 45: 491-497, 1976
 24. Toda N, Hojo M, Sakae K, Usui H: Comparison of the relaxing effect of dopamine with that of adenosine, isoproterenol and acetylcholine in isolated canine coronary arteries. *Blood Vessels* 12: 290-301, 1975
 25. Namm DH, Leader JP: Occurrence and function of cyclic nucleotides in blood vessels. *Blood Vessels* 13: 24-47, 1976
 26. Fog-Møller F, Kemp Genefke I, Bryndum B: Changes in concentration of catecholamines in blood during spontaneous migraine attacks and reserpine-induced attacks. *In* Greene R (ed): *Current Concepts in Migraine Research*. New York, Raven Press, pp 115-119, 1978
 27. Vane JR: Prostacyclin, an endogenous vasodilator and antiplatelet substance generated in the artery wall. *In* Second International Migraine Symposium. The Migraine Trust. London, 1978
 28. Gillilan LA: Blood supply to the central nervous system. *In* Crosby EC, Humphrey T, Lauer EW (eds): *Correlative Anatomy of the Nervous System*. New York, MacMillan Company, pp 550-579, 1972

Local Cerebral Blood Flow in the Conscious Rat As Measured with ¹⁴C-Antipyrine, ¹⁴C-Iodoantipyrine and ³H-Nicotine

K. OHNO, M.D., K. D. PETTIGREW, AND S. I. RAPOPORT, M.D.

SUMMARY Local cerebral blood flow (LCBF) in the conscious rat was estimated with one of 3 radio-tracers — ¹⁴C-antipyrine, ¹⁴C-iodoantipyrine or ³H-nicotine. A tracer was infused intravenously at a constant rate and blood concentration was followed until the animal was killed by decapitation. Tracer concentration was then measured in each of 14 brain regions. The Kety-Schmidt analysis was applied to the data with ¹⁴C-antipyrine and ¹⁴C-iodoantipyrine. The results confirmed findings of Sakurada et al. (1978) that ¹⁴C-iodoantipyrine provides LCBF's that are twice those obtained with ¹⁴C-antipyrine and that approximate LCBF's found with an inert gas. LCBF was calculated from the ³H-nicotine data by assuming complete extraction of tracer from blood by brain. This assumption was approximated for infusion times of 50 sec or less, when LCBF's derived with ³H-nicotine generally did not differ significantly from LCBF's obtained with ¹⁴C-iodoantipyrine. Fifty-sec ³H-nicotine-derived flows for the pineal and pituitary glands were, respectively, 2.59 ± 0.36 (SEM) $cm^3 g^{-1} min^{-1}$ and $1.28 \pm 0.08 cm^3 g^{-1} min^{-1}$. LCBF's calculated for 70 sec to 240 sec of ³H-nicotine infusion were lower than 50-sec values due to back-diffusion, but nevertheless were as high as ¹⁴C-antipyrine LCBF's due to the marked binding of tracer by brain tissue. The results supported the conclusion of Sakurada et al. (1978) that iodoantipyrine is the non-gaseous agent of choice for measuring LCBF precisely, but also showed that short infusion schedules with nicotine provided good estimates of LCBF.

Stroke, Vol 10, No 1, 1979

LOCAL CEREBRAL blood flow (LCBF) can be determined in experimental animals by employing an inert gas tracer, ¹³¹I-trifluoriodomethane, and applying the mathematical principles of Kety to analyze blood-brain exchange.¹⁻⁴ The principles are valid

because diffusional equilibrium between brain and blood is established almost instantaneously.^{5, 6}

Since use of a gas tracer presents technical difficulties, attempts have been made to employ non-gaseous tracers to measure LCBF. ¹⁴C-antipyrine has been used with some success, but it provides LCBF's which are less than those obtained using inert gases. Transfer of ¹⁴C-antipyrine from blood to brain is limited by the comparatively low diffusion coefficient of that tracer in the brain. The cerebrovascular PS value (permeability \times surface area) of ¹⁴C-antipyrine is about $0.020 cm^3 sec^{-1} g^{-1}$, and is not sufficiently high to satisfy the assumptions of the Kety method.^{4, 5, 7-10}

Early work suggested that ¹³¹I-iodoantipyrine might

From the Laboratory of Neurosciences, National Institute on Aging, Gerontology Research Center, Baltimore City Hospital, Baltimore, MD 21224 and Theoretical Statistics and Mathematics Branch, National Institute of Mental Health, Bethesda, MD 20014.

Dr. Ohno is Visiting Fellow, Department of Neurosurgery, Tokyo Medical and Dental University, Tokyo, Japan.

Reprints: Dr. Rapoport, Laboratory of Neurosciences, Gerontology Research Center, Baltimore City Hospital, Baltimore, MD 21224.

be useful for measuring cerebral blood flow,^{11, 12} but it was not until recently that Sakurada et al.⁴ showed that ¹⁴C-iodoantipyrine, which has a higher oil/water partition coefficient than ¹⁴C-antipyrine and also is more permeable at the cerebrovasculature, provides values of LCBF by the Kety method in the conscious rat that are comparable to those obtained with ¹³¹I-trifluoriodomethane. Autoradiography was used to measure local brain concentrations of the ¹⁴C-tracer.

In this paper, we compare LCBF's that were obtained with each of 3 tracers — ¹⁴C-antipyrine, ¹⁴C-iodoantipyrine and ³H-nicotine — in regions of the rat brain that were dissected out according to criteria of Glowinski and Iversen¹³ and Chiueh et al.¹⁴ Knowledge of LCBF in these specific regions is required to calculate regional cerebrovascular permeability to a variety of agents.¹⁵

The mathematical treatment of Kety⁵ was employed to estimate LCBF with both ¹⁴C-antipyrine and ¹⁴C-iodoantipyrine. The treatment could not be used with ³H-nicotine, however, because this tracer binds to brain and is not freely exchangeable between brain and blood.^{10, 16, 17} We estimated LCBF with ³H-nicotine by assuming that all of the tracer was extracted from blood by brain. We chose ³H-nicotine because it, like ¹⁴C-iodoantipyrine, is more permeable than is ¹⁴C-antipyrine at the cerebral vasculature and because its affinity for brain tissue might allow approximately complete extraction at short infusion times.^{10, 17}

Methods

Male rats (Osborn-Mendel strain), weighing 250 g to 350 g, were anesthetized with Na pentobarbital (35 mg kg⁻¹, i.p.). Polyethylene catheters containing 100 IU Na heparin and 0.009 g NaCl/ml of water were implanted in the left femoral artery and vein. The skin was sutured at the catheter exit and infiltrated with 2% butacaine sulfate (Abbott Labs, N. Chicago, Ill.). The hindquarters of the animals were wrapped in a fast-setting plaster bandage (Johnson and Johnson, New Brunswick, N.J.), with hindlimbs and catheters protruding, after which the bandage was tied down on a lead block.⁴ The animals were allowed to recover from anesthesia for 4 hr or more. In their harness, the conscious rats could freely move their forequarters, head and neck, and appeared comfortable.

The catheter in the femoral vein was connected to a 10-ml syringe, which was held in a constant flow pump (Model 255-2, Sage Instruments Inc., White Plains, N.Y.) that was set to deliver fluid at a rate of 0.28 ml min⁻¹. Prior to infusion, a sample of arterial blood was removed for the determination of hematocrit, and for measurement of Pco₂, pH and PaO₂ (pH-Blood Gas Analyzer, No. 213, Instrumentation Labs, Lexington, Mass.). The femoral vein was infused for up to 4 min with isotonic saline containing 20 μC/ml of ³H-nicotine-d-bitartrate (Amersham/Searle, sp. act. = 240 mC/mmol), for 60 sec with isotonic saline containing 4 μC/ml of ¹⁴C-antipyrine (Amersham/Searle, sp. act. = 52 mC/mmol) and for 45 sec with isotonic

saline containing 9 μC/ml of ¹⁴C-iodoantipyrine (New England Nuclear, sp. act. = 50 mC/mmol). The purity of each tracer exceeded 96%, as confirmed by thin layer chromatography. Periodically during infusion, 20 μl samples of arterial blood were collected into heparinized tubes, after which 10 μl aliquots were transferred to scintillation vials.

Animals were decapitated 30 sec to 4 min after infusion began. The skull was opened and the brain was removed and placed on cold filter paper wetted with 0.9% NaCl. Large subarachnoidal and dural blood vessels were removed and discarded. Brain regions then were dissected out, according to the method of Chiueh et al.,¹⁴ and placed in tared scintillation vials that immediately were re-weighed.

The pineal and pituitary glands were removed and the brain was hemisectioned in the midline. The caudate nucleus in the anterior horn of the lateral ventricle, and the hippocampus in the posterior-inferior horn were removed with a curved forceps. The hypothalamus and thalamus then were dissected away from the cerebral cortex and midbrain, using landmarks of the anterior commissure, massa intermedia, mamillary body and internal capsule. The cerebellum subsequently was separated from brain stem, and mid-brain and brain stem were separated at the level below the inferior colliculi. In addition to sampling occipital and frontal cortical and subcortical regions, which contained gray and white matter, gray matter was separated from white matter in the temporal region and corpus callosum. Reproducibility of the dissection procedure has been published.¹⁴

Scintillation vials that contained blood or brain samples received 1.5 ml of Soluene 100 (Packard Instrument Co., Downers Grove, Ill.) and were shaken for 6 hr in a water bath at 60°C. Fifteen ml of Dimilume 30 (Packard Instrument Co.), a liquid scintillation fluid, were added before counting in a Packard Tricarb Liquid Scintillation Spectrometer (Model 2405). Counts per minute (cpm) was converted to disintegrations per minute (dpm), a measure of absolute radioactivity, by using external standardization and predetermined efficiency curves.

Mathematical Analysis of Data

Arterial blood concentrations (C_{blood}dpm/ml) that were measured during i.v. infusion of either ¹⁴C-antipyrine or ¹⁴C-iodoantipyrine were plotted against time and were fit by a non-linear, least-squares procedure with the following equation,¹⁸

$$C_{\text{blood}} = A + Be^{-Rt} + De^{-St} \quad (1)$$

A, B, C, R, and S are constants in Eq. 1 and A + B + C = 0, because C_{blood} = 0 at t = 0 (time when blood concentration starts to rise). Computer fitting provided numerical estimates of the constants from the arterial plasma curve (fig. 1 below).

As shown by Eckman et al.,⁵ a transfer constant K can be defined as follows: where F = LCBF, λ = steady state, tissue: blood partition coefficient for a

particular tracer, and m is a constant between 0 and 1 that represents the extent to which diffusional equilibrium is established between tissue and blood,

$$K = mF/\lambda \quad (2)$$

Local diffusional equilibrium has to be established for the Kety approach to be valid, in which case m approximates 1 and $K = F/\lambda$. F then can be obtained by a least squares fit of the following equation to the data, where T = time of brain sampling and $C_{\text{brain}}(T)$ = tracer concentration (dpm/g) in brain parenchyma (excluding intravascular concentration) at time T , and only K is unknown,

$$C_{\text{brain}}(T)/\lambda = A - (A + BK/[K - R] + DK/[K - S])e^{-KT} + (K/[K - R])Be^{-RT} + (K/[K - S])De^{-ST} \quad (3)$$

K was calculated with Eq. 3 by a non-linear iterative least squares process that employed the MLAB program on a PDP-10 computer.^{17, 18} F was obtained from K by Eq. 2, letting $m = 1$ and $\lambda = 0.9$ for ^{14}C -antipyrine and 0.8 for ^{14}C -iodoantipyrine.⁴

Results

Figure 1 illustrates blood concentrations of ^{14}C -iodoantipyrine during 45 sec of infusion, as well as the least squares fit of Eq. 1 to these data. The constants that were derived by the fit were inserted into Eq. 3, together with $m = 1$, $\lambda = 0.8^4$ and $C_{\text{brain}}(T)$. $C_{\text{brain}}(T)$, which represents intraparenchymal brain concentration of tracer, was obtained by subtracting intravascular from net regional radioactivity when the former quantity was taken as the product of regional blood volume and blood concentration (dpm/ml). Regional volumes are as follows (% of wet wt): olfactory bulb (4.68), caudate nucleus (1.24), hippocampus (1.61), frontal lobe (2.09), occipital lobe (2.42), thalamus + hypothalamus (1.79), colliculi (2.01), cerebellum (3.46), pons (2.56), medulla (3.53), gray matter (parietal) (2.70), white matter (corpus callosum) (1.14). Regional blood volume at the pineal and pituitary glands was taken as 5% of wet wt.^{16, 19}

Mean hematocrit, arterial pH, PaO_2 and PaCO_2 did not differ significantly between experimental groups ($P > 0.05$). For 7 animals infused either with ^{14}C -antipyrine or ^{14}C -iodoantipyrine, means \pm SEM's were as follows: hematocrit = $48.0 \pm 0.24\%$, $\text{PaCO}_2 = 38.4 \pm 1.8$ mm Hg, $\text{PaO}_2 = 83.8 \pm 3.2$ mm Hg and pH = 7.41 ± 0.013 units. For 10 animals administered ^3H -nicotine, means \pm SEM's were: hematocrit = $47.4 \pm 0.81\%$, $\text{PaCO}_2 = 39.2 \pm 0.46$ mm Hg, $\text{PaO}_2 = 84.1 \pm 0.81$ mm Hg and pH = 7.41 ± 0.02 units.

Table 1 lists mean LCBF's that were calculated by Eqs. 1 and 3 from data obtained with ^{14}C -antipyrine and ^{14}C -iodoantipyrine. The results confirm the original findings of Sakurada et al.⁴ that ^{14}C -iodoantipyrine provides about 2-fold higher values of LCBF than does ^{14}C -antipyrine. Furthermore, the LCBF's in




FIGURE 1. Arterial blood ^{14}C concentration during i.v. infusion of ^{14}C -iodoantipyrine in the conscious rat. Note that t in the equation equals Infusion Time — 8 sec.

table 1 generally do not differ significantly from published values by Sakurada et al.,⁴ although LCBF's for ^{14}C -antipyrine generally are less than ^{14}C -antipyrine LCBF's reported by Ginsberg et al.²⁰ As compared to the mean values in table 1, LCBF's reported by Sakurada et al.,⁴ are larger at the cerebellum with ^{14}C -antipyrine, and smaller at white matter with both ^{14}C -antipyrine and ^{14}C -iodoantipyrine. Not all their data are comparable to data in table 1, however, as many regions in the table include several areas that were analyzed separately by their more discriminating autoradiographic method.

TABLE 1 LCBF Calculated with ^{14}C -Antipyrine and ^{14}C -Iodoantipyrine

Structure	Local cerebral blood flow, $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$	
	^{14}C -Antipyrine	^{14}C -Iodoantipyrine
Olfactory bulb	$0.59 \pm 0.03(6)^{a,b}$	$1.02 \pm 0.10(6)$
Caudate nucleus	0.62 ± 0.04^b	1.80 ± 0.19
Hippocampus	0.60 ± 0.05^b	1.38 ± 0.16
Frontal lobe	0.73 ± 0.05^b	1.68 ± 0.29
Occipital lobe	0.70 ± 0.06^b	1.80 ± 0.29
Thalamus + hypothalamus	0.62 ± 0.05^b	1.50 ± 0.17
Superior colliculus	0.67 ± 0.05^b	1.68 ± 0.17
Inferior colliculus	0.80 ± 0.08^b	2.04 ± 0.25
Cerebellum	0.58 ± 0.04^b	1.02 ± 0.11
Pons	0.56 ± 0.04^b	1.26 ± 0.12
Medulla	0.49 ± 0.05^b	1.20 ± 0.13
Gray matter, parietal	0.86 ± 0.08^b	2.40 ± 0.37
White matter, corpus callosum	0.43 ± 0.04^b	1.14 ± 0.14
Pineal gland	0.61 ± 0.40	—
Pituitary gland	0.74 ± 0.16	1.04 ± 0.15

^aMean \pm SEM (No. of experiments in column is in parenthesis).

^bDiffers significantly from ^{14}C -iodoantipyrine mean ($p < 0.05$).

The Kety analysis, which assumes diffusional equilibrium between plasma and brain, cannot be employed if the tracer is bound or incorporated within the brain as is ³H-nicotine.^{10, 17} We therefore estimated LCBF with ³H-nicotine when assuming complete extraction by brain, and evaluated this assumption at different durations of i.v. infusion. LCBF was calculated as the ratio of net measured regional brain ³H content at the time of death (T), dpm/g [parenchymal tracer C_{brain}(T) plus intravascular tracer (C_{blood}(T) × Blood Volume)], divided by the integrated blood concentration curve up to this time,

$$LCBF = \frac{\text{Net brain concentration (T)}}{\int_0^T C_{\text{blood}} dt} \quad (4)$$

Table 2 presents mean LCBF's that were calculated by Eq. 4 at infusion times of 30 to 240 sec, and figure 2 presents these means when normalized to the maximal mean for a specific region, taken as 100% and usually at 50 sec of infusion. In no instance did the 30-sec mean differ significantly from the 50-sec mean LCBF, although LCBF's derived at infusion times of 70 sec or more generally were significantly less than respective 50-sec values (*P* < 0.05). With some few exceptions (cf. table 1), 30-sec and 50-sec LCBF's derived with ³H-nicotine did not differ from LCBF's derived with ¹⁴C-iodoantipyrine but did exceed ¹⁴C-antipyrine LCBF's.

As illustrated by figure 3, binding of ³H-nicotine by brain was markedly evident at infusion times of 70 sec to 240 sec, when brain concentration of tracer exceeded blood concentration by as much as 4-fold. For




FIGURE 2. LCBF obtained with ³H-nicotine in rats infused intravenously for between 30 and 240 sec, given as percent of maximal value for each brain region.

infusion times of 50 sec or less, blood concentration usually exceeded brain concentration. This inequality and the large binding of intracerebral tracer reduced back-diffusion from brain to such an extent that Eq. 4 could be used to afford LCBF's that approximated those obtained with ¹⁴C-iodoantipyrine.

Discussion

These findings on dissected brain regions confirm conclusions obtained by autoradiographic procedures that ¹⁴C-iodoantipyrine provides higher values of LCBF than does ¹⁴C-antipyrine in the conscious rat.⁴

TABLE 2 Calculated LCBF with ³H-nicotine in Relation to i.v. Infusion Time. Mean LCBF's at Infusion Times of 30, 70, 120, 240 sec were Compared to LCBF's at 50 sec by a Multiple Comparisons Procedure, Using Bonferroni *t* Statistics.³⁸

Structure	Infusion time (sec)				
	30	50	70	120	240
	Local cerebral blood flow, cm ³ g ⁻¹ min ⁻¹				
Olfactory bulb	1.11 ± 0.12(6) ^{a,c}	1.23 ± 0.08(10) ^c	0.87 ± 0.06(6) ^b	0.78 ± 0.08(3) ^b	0.73 ± 0.07(3) ^b
Caudate nucleus	1.17 ± 0.11 ^{c,d}	1.28 ± 0.08 ^{c,d}	1.01 ± 0.10 ^c	0.77 ± 0.19 ^b	0.81 ± 0.08 ^b
Hippocampus	1.04 ± 0.10 ^c	1.14 ± 0.08 ^c	0.84 ± 0.07	0.85 ± 0.11	0.71 ± 0.06 ^b
Frontal lobe	1.74 ± 0.16 ^c	1.82 ± 0.17 ^c	1.23 ± 0.15	1.04 ± 0.28	0.95 ± 0.07 ^b
Occipital lobe	1.57 ± 0.15 ^c	1.80 ± 0.12 ^c	1.44 ± 0.13 ^c	0.85 ± 0.17 ^b	0.84 ± 0.07 ^b
Thal. + hypothal.	1.03 ± 0.08 ^{c,d}	1.23 ± 0.07 ^c	0.87 ± 0.06 ^b	0.83 ± 0.15 ^b	0.77 ± 0.04 ^b
Sup. colliculus	1.08 ± 0.12 ^{c,d}	1.26 ± 0.08 ^{c,d}	0.86 ± 0.05 ^b	1.04 ± 0.16	0.78 ± 0.04 ^b
Inf. colliculus	1.28 ± 0.16 ^c	1.36 ± 0.07 ^c	1.03 ± 0.06	1.03 ± 0.11	0.80 ± 0.04 ^b
Cerebellum	0.79 ± 0.09	0.91 ± 0.04 ^c	0.71 ± 0.04 ^b	0.64 ± 0.08 ^b	0.58 ± 0.05 ^b
Pons	0.95 ± 0.07 ^c	1.11 ± 0.07 ^c	0.78 ± 0.05 ^b	0.80 ± 0.07 ^b	0.62 ± 0.03 ^b
Medulla	0.85 ± 0.09 ^c	1.10 ± 0.06 ^c	0.75 ± 0.05 ^b	0.81 ± 0.05 ^c	0.62 ± 0.04 ^b
Gray matter	2.38 ± 0.19 ^c	2.81 ± 0.21 ^c	1.72 ± 0.16 ^b	1.23 ± 0.37 ^b	1.08 ± 0.03 ^b
White matter	0.59 ± 0.05 ^d	0.73 ± 0.06 ^{c,d}	0.53 ± 0.05 ^b	0.44 ± 0.12 ^b	0.51 ± 0.03
Pineal gland	4.16 ± 1.54	2.59 ± 0.36	1.63 ± 0.53	1.91 ± 1.04	1.57 ± 0.21
Pituitary gland	1.21 ± 0.24	1.28 ± 0.08	0.96 ± 0.08	1.03 ± 0.19	0.69 ± 0.15

^aMean ± SEM (No. of experiments in column is in parenthesis); ^bDiffers significantly from 50-sec ³H-nicotine mean (*p* < 0.05); ^cDiffers significantly from ¹⁴C-antipyrine mean in table 1 (*p* < 0.05); ^dDiffers significantly from ¹⁴C-iodoantipyrine mean in table 1 (*p* < 0.05), where only 30-sec and 50-sec ³H-nicotine means are considered.




FIGURE 3. ^3H concentration (dpm/ml) in whole blood of rats during *i.v.* infusion of ^3H -nicotine for 30 sec to 240 sec, as well as regional brain parenchymal concentrations, C_{brain} dpm/g, at time of death in individual experiments. Concentrations were normalized to the 30-sec blood concentration (taken as 1).

^{14}C -iodoantipyrine also furnishes higher LCBF's than does the hydrogen clearance method. For example, at the frontal cortex of the conscious rat, LCBF equals 0.79 ± 0.22 (SEM) $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ with the hydrogen clearance method,²¹ but equals 1.68 ± 0.29 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ with ^{14}C -iodoantipyrine (table 1).

The Kety method^{1, 2} was designed for use with an inert gas, such as ^{131}I -trifluoriodomethane, that can reach almost instantaneous equilibrium between brain and blood. This condition is not satisfied by ^{14}C -antipyrine, which gives low LCBF's, because its entry into brain is somewhat limited by diffusion at the capillary and does not depend only on blood flow.^{4-6, 8, 17} Diffusional equilibrium presumably is reached, on the other hand, by the more lipid-soluble ^{14}C -iodoantipyrine, which provides LCBF's that compare with those obtained with ^{131}I -trifluoriodomethane.⁴ ^{14}C -iodoantipyrine would appear to be the non-gaseous agent of choice for exact measurement of LCBF, not only because it gives values comparable to those with

the inert gas, but also because it is long-lived within blood, not significantly metabolized within 1 min, commercially available and exhibits a tissue-blood partition coefficient that is uniform throughout the brain and is independent of hematocrit.⁴

Due to its possible back-diffusion from brain to blood, ^3H -nicotine cannot be used with certainty to obtain LCBF. However, the equivalent values of LCBF as derived with 30 or 50 sec of infusion of ^3H -nicotine, as compared to those provided by ^{14}C -iodoantipyrine and the Kety analysis, demonstrate that back-diffusion was minimal for infusion times of up to 50 sec, and that Eq. 4 then can be used to derive LCBF precisely. The lower LCBF's obtained with ^3H -nicotine than with ^{14}C -iodoantipyrine in occasional regions (thalamus + hypothalamus at 30 sec of infusion; superior colliculus, caudate nucleus and white matter at 30 and 50 sec), may have been due to incomplete extraction of ^3H -nicotine by those regions or to some back-diffusion into blood.

For 120 sec or 240 sec of ^3H -nicotine infusion, delivery rate may be less important than tissue uptake in calculating LCBF. The higher calculated LCBF of gray as compared to white matter reflects in part specific, reversible uptake of ^3H -nicotine by cell bodies and nerve endings, in part at cholinergic receptors. Uptake is lowered *in vivo* by pentobarbital anesthesia, and is reduced in brain slices by lowering temperature.^{16, 22-27} Most of the ^3H in the brain at 240 sec is associated with ^3H -nicotine; less than 3% is due to ^3H -cotinine, a metabolic derivative of ^3H -nicotine that is produced in liver and kidney and appears in blood about 180 sec after an intravenous injection, but is poorly permeable at the blood-brain barrier.^{26, 28}

The pineal and pituitary glands are outside of the blood-brain barrier system.²⁹ Local blood flows in these glands, as measured with ^{14}C -iodoantipyrine and ^3H -nicotine, are comparable to flows derived by analyzing the distribution of cardiac output to these glands — 2 to 3.43 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ at the pineal gland, 1.01 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ at the anterior pituitary and 3.74 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ at the posterior pituitary.^{19, 30, 31} ^{125}I -antipyrine gives flows of 1.2 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ at the anterior pituitary and 2.36 $\text{cm}^3\text{g}^{-1}\text{min}^{-1}$ at the posterior pituitary of the conscious rat.³²

LCBF closely correlates with regional cerebral metabolism under many physiological conditions in the same animal, and when comparing brains of different species.³³ Both LCBF and cerebral energy metabolism are higher, for example, in the rat than in many other species, possibly because cellular packing is greater in the rat brain.^{4, 34-36} If ^3H -nicotine binding by brain is a measure of density of cells and nerve endings, then short-term experiments (50 sec or less) with nicotine might be used to estimate LCBF and long-term experiments (5 min or more) to estimate density of cells and nerve endings.^{16, 22-27} Nicotine, when incorporating the positron emitter ^{13}N , might also be used in man to measure regional blood flow by means of emission computed tomography.³⁷

Acknowledgment

We thank Drs. O. Sakurada and L. Sokoloff for providing us with a preprint of their publication on LCBF, and for their helpful suggestions and criticism.

References

- Kety SS: The theory and applications of the exchange of inert gas at the lungs and tissues. *Pharmacol Rev* 3: 1-41, 1951
- Kety SS: Measurement of local blood flow by the exchange of an inert, diffusible substance. *Methods Med Res* 8: 228-236, 1960
- Freygang WH Jr, Sokoloff L: Quantitative measurement of regional circulation in the central nervous system by the use of radioactive inert gas. *Adv Biol Med Phys* 6: 263-279, 1958
- Sakurada O, Kennedy C, Jehle J, Brown JD, Carbin GL, Sokoloff L: Measurement of local cerebral blood flow with iodo [¹⁴C] antipyrine. *Am J Physiol* 234(1): H59-H66, 1978
- Eckman WW, Phair RD, Fenstermacher JD, Patlak CS, Kennedy C, Sokoloff L: Permeability limitation in estimation of local brain blood flow with [¹⁴C] antipyrine. *Am J Physiol* 229: 215-221, 1975
- McHenry LC Jr, Jaffe ME, Goldberg HI: Regional cerebral blood flow measurement with small probes. I. Evaluation of the method. *Neurology (Minneapolis)* 19: 1198-1206, 1969
- Crone C: The permeability of brain capillaries to nonelectrolytes. *Acta Physiol Scand* 64: 407-417, 1965
- Eklof B, Lassen NA, Nilsson L, Norberg K, Siesjö BK, Torlöf P: Regional cerebral blood flow in the rat measured by the tissue sampling technique; a critical evaluation using four indicators C¹⁴-antipyrine, C¹⁴-ethanol, H³-water and Xenon¹³³. *Acta Physiol Scand* 91: 1-10, 1974
- Oldendorf WH: Lipid solubility and drug penetration of the blood-brain barrier. *Proc Soc Exp Biol Med* 147: 813-816, 1974
- Oldendorf WH: Brain concentrations of ¹⁴C-nicotine and ¹⁴C-antipyrine after intravenous injection. pp 103-109. *In* Langfitt TN, McHenry LC, Reivich M, Wallmann H (eds) *Cerebral Circulation and Metabolism*. New York, Springer-Verlag 1975
- Sapirstein LA: Regional blood flow by fractional distribution of indicators. *Am J Physiol* 193: 161-168, 1958
- Reinmuth OM, Scheinberg P, Bourne B: Total cerebral blood flow and metabolism. *Arch Neurol* 12: 49-66, 1965
- Głowinski J, Iversen LL: Regional studies of catecholamines in the rat brain. I. The disposition of [³H] norepinephrine, [³H] dopamine and [³H] DOPA in various regions of the brain. *J Neurochem* 13: 655-669, 1966
- Chiueh CC, Sun CL, Kopin IJ, Fredericks WR, Rapoport SI: Entry of ³H-norepinephrine, ¹²⁵I-albumin and Evans blue from blood into brain following unilateral osmotic opening of the blood-brain barrier. *Brain Res* 145: 291-301, 1978
- Ohno K, Pettigrew KD, Rapoport SI: Lower limits of cerebrovascular permeability to nonelectrolytes in the conscious rat. *Am J Physiol* 253 (3), H299-H307, 1978
- Schmitterlöw RC, Hansson E, Andersson G, Applegren L-E, Hoffmann PC: Distribution of nicotine in the central nervous system. *Ann NY Acad Sci* 142: 2-14, 1967
- Bradbury MWB, Patlak CS, Oldendorf WH: Analysis of brain uptake and loss of radiotracers after intracarotid injection. *Am J Physiol* 229: 1110-1115, 1975
- Knott GD, Shrager RI: On-line modeling by curve fitting. pp 138-151. *In* *Computer Graphics: Proceedings of the SIGGRAPH Computers in Medicine Symposium 6* (no 4). ACM, SIGGRAPH Notices, 1972
- Goldman H, Wurtman RJ: Flow of blood to the pineal body of the rat. *Nature* 203: 87-88, 1964
- Ginsberg MD, Medoff R, Reivich M: Heterogeneities of regional cerebral blood flow during hypoxia-ischemia in the rat. *Stroke* 7: 132-134, 1976
- Haining JL, Turner MD, Pantall RM: Measurement of local cerebral blood flow in the unanesthetized rat using a hydrogen clearance method. *Circ Res* 23: 313-324, 1968
- Alderdice MT, Weiss GB: Effects of pharmacological agents on [¹⁴C]-nicotine distribution and movements in slices from different rat brain areas. *Neuropharmacology* 14: 811-817, 1975
- Weiss GB, Alderdice MT: Characterization of [¹⁴C]-nicotine accumulation and movements in slices from different brain areas. *Neuropharmacology* 14: 265-273, 1975
- Tsujimoto A, Nakashima T, Tanino S, Dohi T, Kuroguchi Y: Tissue distribution of [³H] nicotine in dogs and rhesus monkeys. *Toxicol Appl Pharmacol* 32: 21-31, 1975
- Eterović VA, Bennett EL: Nicotinic cholinergic receptor in brain detected by binding of α -[³H] bungarotoxin. *Biochim Biophys Acta* 362: 346-355, 1974
- Appelgren L-E, Hansson E, Schmitterlöw CG: The accumulation and metabolism of C¹⁴-labelled nicotine in the brain of mice and cats. *Acta Physiol Scand* 56: 249-257, 1962
- Brown DA, Halliwell JV: Intracellular pH in rat isolated superior cervical ganglia in relation to nicotine-depolarization and nicotine-uptake. *Br J Pharmacol* 45: 349-359, 1972
- Turner DM: Metabolism of small multiple doses of [¹⁴C]-nicotine in the cat. *Br J Pharm* 41: 521-529, 1971
- Rapoport SI: *Blood-Brain Barrier in Physiology and Medicine*. New York, Raven Press, 1976
- Goldman H: The nervous control of blood flow to the pineal body. *Life Sci* 6: 2071-2077, 1967
- Goldman H: Failure of cervical sympathectomy to alter pituitary blood flow. *Endocrinology* 83: 603-606, 1968
- Lichardus B, Albrecht I, Ponc J, Linhart L: Water deprivation for 24 hours increases selectively blood flow in posterior pituitary of conscious rats. *Endocrinol Exp (British)* 11: 99-104, 1977
- Reivich M, Sokoloff L: Application of the 2-deoxy-D-glucose method to the coupling of cerebral metabolism and blood flow. *Neurosci Res Program Bull* 14: 474-475, 1976
- Landau WM, Freygang WH, Roland LP, Sokoloff, Kety SS: The local circulation of the living brain; values in the unanesthetized and anesthetized cat. *Trans Am Neurol Assoc* 80: 125-129, 1955
- Nilsson B, Siesjö BK: A method for determining blood flow and oxygen consumption in the rat brain. *Acta Physiol Scand* 96: 72-82, 1976
- Sokoloff L, Reivich M, Kennedy C, Des Rosiers MH, Patlak CS, Pettigrew KD, Sakurada O, Shinohara M: The [¹⁴C] deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *J Neurochem* 28: 897-916, 1977
- Phelps ME: Emission computed tomography. *Seminars in Nuclear Medicine* 7: 337-365, 1977
- Miller RG: *Simultaneous Statistical Inference*. pp 67-70. New York, McGraw-Hill

Local cerebral blood flow in the conscious rat as measured with ¹⁴C-antipyrine, ¹⁴C-iodoantipyrine and ³H-nicotine.
K Ohno, K D Pettigrew and S I Rapoport

Stroke. 1979;10:62-67

doi: 10.1161/01.STR.10.1.62

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 1979 American Heart Association, Inc. All rights reserved.

Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://stroke.ahajournals.org/content/10/1/62.citation>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Stroke* is online at:
<http://stroke.ahajournals.org/subscriptions/>