Low Pessimism Protects Against Stroke The Health and Social Support (HeSSup) Prospective Cohort Study

Hermann Nabi, PhD; Markku Koskenvuo, MD, PhD; Archana Singh-Manoux, PhD; Jyrki Korkeila, MD, PhD; Sakari Suominen, MD, PhD; Katariina Korkeila, MD, PhD; Jussi Vahtera, MD, PhD; Mika Kivimäki, PhD

- **Background and Purpose**—The association between optimism and pessimism and health outcomes has attracted increasing research interest. To date, the association between these psychological variables and risk of stroke remains unclear. We examined the relationship between pessimism and the 7-year incidence of stroke.
- *Methods*—A random sample of 23 216 adults (9480 men, 13 796 women) aged 20 to 54 years completed the pessimism scale in 1998, that is, at study baseline. Fatal and first nonfatal stroke events during a mean follow-up of 7.0 years were documented by linkage to the national hospital discharge and mortality registers leading to 105 events.
- *Results*—Unadjusted hazard ratio was 0.44 (95% CI, 0.25 to 0.77) for participants in the lowest quartile (a low pessimism level) when compared with those in the highest quartile (a high pessimism level). After serial adjustments for sociodemographic characteristics, cardiovascular biobehavioral risk factors, depression, general feeling of stressfulness, and ischemic heart disease, the fully adjusted hazard ratio was 0.52 (95% CI, 0.29 to 0.93).

Conclusions—In this population of adult men and women, low level of pessimism had a robust association with reduced incidence of stroke. (*Stroke*. 2010;41:187-190.)

Key Words: epidemiology ■ pessimism ■ psych & ■ behavior

There is a growing research interest in the relationship between dispositional optimism and pessimism, defined as a general tendency to exhibit positive and negative expectancies about the future, and health outcomes.^{1–3} Although findings from well-conducted prospective studies suggest an association between optimism and pessimism as separate constructs and the risk of incident coronary heart disease (CHD) and cardiovascular disease mortality,^{1,4,5} no evidence to date is available of an association with the incidence of stroke. We used prospective data from a large sample of the Finnish population to examine the association between dispositional pessimism and the incidence of stroke.

Methods

Population

The Health and Social Support (HeSSup) study is a prospective cohort study on a population sample representative of the Finnish population of the following 4 age groups: 20 to 24, 30 to 34, 40 to 44, and 50 to 54 years at baseline in 1998; there was a total of 10 628 men and 15 267 women. The Turku University Central Hospital Ethics Committee approved the study.

Dispositional pessimism was assessed using the revised Life Orientation Test–Revised (LOT-R)^{6,7} (Cronbach's α =0.74). The measure includes 6 statements, of which 3 are worded positively for optimism and 3 are worded negatively to indicate pessimism. We categorized pessimism mean scores into 4 groups based on the nearest approximate of the quartiles as in previous studies (Supplemental Table I, available online at http://stroke.ahajournals.org).⁴ The highest quartile (high pessimism) was the reference category in the analysis.

Follow-Up of Stroke Events

Linkage to the National Hospital Discharge Register and the Statistics Finland Mortality Register provided a virtually complete follow-up for hospitalizations and death for all participants who were treated in a hospital or died between January 1, 1999, and December 31, 2005. Stroke was determined by the International Classification of Diseases, 10th Edition codes I60 (subarachnoid hemorrhage), I61 (intracerebral hemorrhage), and I63 (cerebral infarction) as the main diagnosis of hospitalization or death.

Cardiovascular Risk Factors

History of hypertension, coronary heart disease, and diabetes mellitus at the time of the questionnaire in 1998 (ie, the survey year) was ascertained based on strict criteria.⁸ We excluded all participants hospitalized for CHD or cardiovascular disease and those with

© 2009 American Heart Association, Inc.

Stroke is available at http://stroke.ahajournals.org

From INSERM U687-IFR69 (H.N., A.S.-M.), Villejuif, France; the Department of Public Health (M. Koshenvuo), University of Helsinki, Helsinki, Finland; the Department of Epidemiology and Public Health (A.S.-M., M. Kivimäki), University College London, London, UK; the Department of Psychiatry (J.K.), University of Turku and Harjavalta Hospital, Turku, Finland; the Department of Public Health (S.S., J.V.), University of Turku and Turku University Hospital, Turku, Finland; Municipal Health Care (K.K.), Turku, Finland; and the Finnish Institute of Occupational Health (J.V., M. Kivimäki), Helsinki, Finland.

Correspondence to Hermann Nabi, PhD, INSERM Unité 687, 16 avenue Paul Vaillant Couturier, 94807 Villejuif Cedex, France. E-mail Hermann.Nabi@inserm.fr

Table 1.	Sample	Characteristics	as a	Function	of the N	lo. of	Stroke	Events	and	Pessimism	Score	Levels
----------	--------	-----------------	------	----------	----------	--------	--------	--------	-----	-----------	-------	--------

Baselin Covariates	No. Events/No. Participants	1 (Lowest)	I (Lowest) 2		4 (Highest)	P Value for Trend	
Sex						0.243	
Men	59/9480	3124 (33.0)	2333 (24.6)	2268 (23.9)	1755 (18.5)		
Women	46/13736	4467 (32.5)	3571 (26.0)	3319 (24.2)	2379 (17.3)		
Age group, years						0.001	
20–24	3/6324	1938 (30.6)	1518 (24.0)	1664 (26.3)	1204 (19.0)		
30–34	8/5533	1937 (35.0)	1399 (25.3)	1273 (23.0)	924 (16.7)		
40–44	28/5555	1831 (33.0)	1473 (26.5)	1262 (22.7)	989 (17.8)		
50–54	66/5804	1885 (32.5)	1514 (26.1)	1388 (23.9)	1017 (17.5)		
Level of education						< 0.001	
Basic (compulsory)	44/7412	2136 (28.8)	1756 (23.7)	1911 (25.8)	1609 (21.7)		
Secondary	24/5232	1367 (26.1)	1273 (24.3)	1441 (27.5)	1151 (22.0)		
College	26/7400	2696 (36.4)	2033 (27.5)	1654 (22.4)	1017 (13.7)		
University	11/3172	1392 (43.9)	842 (26.5)	581 (18.3)	357 (11.3)		
Marital status						< 0.001	
Other	25/7674	2325 (30.3)	1735 (22.6)	1881 (24.5)	1733 (22.6)		
Married/cohabiting	80/15542	5266 (33.9)	4169 (26.8)	3706 (23.8)	2401 (15.4)		
Current smoker						< 0.001	
No	62/15508	5256 (33.9)	4114 (26.5)	3714 (23.9)	2424 (15.6)		
Yes	35/5862	1717 (29.3)	1311 (22.4)	1441 (24.6)	1393 (23.8)		
Missing	8/1846	618 (33.5)	479 (25.9)	432 (23.4)	317 (17.2)		
High alcohol intake (200 g of alcohol/week*)						< 0.001	
No	87/20956	6950 (33.2)	5412 (25.8)	5034 (24.0)	3560 (17.0)		
Yes	18/2260	641 (28.4)	492 (21.8)	553 (24.5)	574 (25.4)		
Obesity (body mass index \geq 30 kg/m ²)						< 0.001	
No	83/20992	6973 (33.2)	5395 (25.7)	5050 (24.1)	3574 (17.0)		
Yes	22/2224	618 (27.8)	509 (22.9)	537 (24.1)	560 (25.2)		
Sedentary lifestyle (<2 MET hours per day)						< 0.001	
No	69/17855	6072 (34.0)	4672 (26.2)	4205 (23.6)	2906 (16.3)		
Yes	36/5361	1519 (28.3)	1232 (23.0)	1382 (25.8)	1228 (22.9)		
Depression (Beck Depression Inventory score \geq 10)						< 0.001	
No	97/22159	7559 (34.1)	5833 (26.3)	5408 (24.4)	3359 (15.2)		
Yes	8/1057	32 (3.0)	71 (6.7)	179 (16.9)	775 (73.3)		
General feeling of stressfulness in daily life						0.715	
Low	46/10497	3464 (33.0)	2604 (24.8)	2574 (24.5)	1855 (17.7)		
Intermediate	27/5547	1790 (32.3)	1426 (25.7)	1312 (23.7)	1019 (18.4)		
High	31/7020	2294 (32.7)	1833 (26.1)	1666 (23.7)	1227 (17.5)		
Missing	1/152	43 (28.3)	41 (27.0)	35 (23.0)	33 (21.7)		
Hypertension or diabetes						< 0.001	
No	85/22136	7299 (33.0)	5648 (25.5)	5300 (23.9)	3889 (17.6)		
Yes	20/1080	292 (27.0)	256 (23.7)	287 (26.6)	245 (22.7)		
Ischemic heart disease						0.002	
No	102/23016	7566 (32.7)	5864 (25.5)	5533 (24.0)	4083 (17.7)		
Yes	3/200	55 (27.5)	40 (20.0)	54 (27.0)	51 (25.5)		

*A total of 200 g of alcohol represent 16 drinks per week.

MET indicates metabolic equivalent task.

medication for CHD during or before 1998. Incident CHD events (International Classification of Diseases, 10th Edition codes I20 to I25) that occurred during the follow-up were used as a covariate in the analysis. Behavior-related risk factors include current smoking status, alcohol consumption, physical activity, and obesity. Depressive symptoms were assessed using the Beck Depression Inventory

(score ${<}10$ versus 10+). The general feeling of stressfulness in daily life was measured using the Reeder Stress Inventory.⁹

Statistical Analysis

We examined the relationship between pessimism and subsequent stroke events using 7 serially adjusted Cox regression models. We

Table 2	. Hazard Ratio	os (95% Cls) for th	e Association Bet	tween Quartiles of	Pessimism Score	and Incident Stroke Events
---------	----------------	---------------------	-------------------	--------------------	-----------------	----------------------------

	Quartiles of Pessimism Score						
Adjustments	1 (Lowest; 22 Events/n=7588)	2 (27/5904)	3 (29/5587)	4 (Highest; 27/4127)			
Model 1: none	0.44 (0.25-0.77)†	0.69 (0.41–1.18)	0.79 (0.46-1.34)	1			
Model 2: sex, age, education, marital status	0.47 (0.26–0.84)†	0.71 (0.42–1.23)	0.82 (0.48–1.39)	1			
Model 3: sex, age, education, marital status+behavior-related risk factors	0.51 (0.25-0.91)*	0.79 (0.46–1.36)	0.89 (0.52-1.49)	1			
Model 4: sex, age, education, marital status+hypertension, and diabetes	0.48 (0.27–0.86)†	0.73 (0.43–1.26)	0.83 (0.49–1.39)	1			
Model 5: sex, age, education, marital status+depression	0.48 (0.26–0.87)†	0.73 (0.41–1.28)	0.83 (0.48–1.44)	1			
Model 6: sex, age, education, marital status+general feeling of stressfulness	0.47 (0.270.84)†	0.72 (0.42–1.24)	0.82 (0.49–1.39)	1			
Model 7: sex, age, education, marital status+incident CHD	0.44 (0.25–0.77)†	0.68 (0.40–1.16)	0.80 (0.47-1.36)	1			
Model 8: all aforementioned	0.52 (0.29–0.93)*	0.80 (0.47–1.39)	0.88 (0.52–1.50)	1			

**P*<0.05.

†P<0.01.

Behavior-related risk factors are smoking, alcohol consumption, sedentary lifestyle, obesity.

combined men and women in the analyses (P>0.05 for interaction with sex) and verified that the assumption of proportional hazards held (all P>0.05).

Results

Data linkage to national health registers, on the basis of a written consent, was successful for 24 128 (93%) participants. A total of 23 216 participants had complete data on the pessimism scale and all covariates. Of them, 105 incident stroke events (fatal and nonfatal 25 subarachnoid hemorrhages, 23 intracerebral hemorrhages, and 57 cerebral infarctions) were documented during a mean follow-up of 7.0 years (range, 0.02 to 7.01 years; Table 1).

Table 2 presents the association of pessimism levels with incident stroke events. Compared with participants with a high level of pessimism (the highest quartile), those reporting a low level of pessimism (the lowest quartile) were at lower risk of stroke (hazard ratio=0.44, 95% CI=0.25 to 0.77). After serial adjustments, the fully adjusted hazard for stroke was 0.52 (95% CI=0.29 to 0.93) among those with low pessimism.

The survival curve for stroke among participants with the lowest quartile of pessimism differed significantly (log rank test, P < 0.05) from those in the other quartiles and the difference between the groups grew as a function of time (Figure).

Further analyses were run examining the role of optimism (in quartiles) in the risk of stroke. We found no evidence to suggest an association between optimism and the incidence of stroke (P=0.53), lending support to the idea that pessimism and optimism are separate constructs. We also examined the association between pessimism and stroke incidence in analysis adjusted for optimism in addition to age, sex, education, and marital status. The adjustment did not alter the reduced incidence of stroke associated with low pessimism (hazard ratio=0.49, 95% CI=0.26 to 0.90).

Discussion

We examined the prospective association between dispositional pessimism and subsequent stroke events in a large population of men and women with no history of ischemic heart disease or cardiovascular disease. We found that participants with a low pessimism level had a 48% lower risk for incident stroke over the 7 years of follow-up after controlling for many potential confounders. We found no evidence to suggest that the association was attributable to depressive symptoms, a correlate of both pessimism and the risk of stroke,^{10,11} It must be noted that we assessed depressive symptoms with the Beck Depression Inventory, which is a validated survey instrument but not a diagnostic tool for clinical depression.

The present study was based on a representative sample of the Finnish population in 4 age groups but did not include elderly participants, which may limit the generalizability of our findings. Our findings are broadly consistent with previous studies on the associations between pessimism and optimism and the risk of CHD or death,^{4,12} Our results highlight the importance of pessimism in particular for the incidence of stroke, even in analyses adjusted for optimism. In agreement with the present results, a recent clinical study showed a lower survival rate for pessimistic patients with



Figure. Cumulative incidence of stroke (fully adjusted*) by pessimism score quartiles. *Sex, age, education, marital status, smoking, alcohol consumption, sedentary lifestyle, obesity, hypertension and diabetes, depression, general feeling of stressfulness, and incident CHD.

cancer than their less pessimistic counterparts but no difference between individuals with high levels of optimism and those with low levels of optimism.¹³ The precise mechanisms underlying the link between pessimism and stroke remain unclear. Both behavioral (lifestyle behaviors) and biological (autonomic nervous system activity) mechanisms are plausible.^{5,10,11,14} Low pessimism may also be related to favorable trajectories of stroke risk factors over time. Further longitudinal studies are needed to examine these mechanisms in detail and whether interventions designed to reduce pessimism would alter stroke risk.

Acknowledgments

The HeSSup study is supported by the Academy of Finland (3 grants), the Yrjö Jahnson Foundation (3 grants), and the Finnish Heart Foundation (one grant). M. Kivimäki and J.V. are supported by the Academy of Finland (grants 117604, 124271, 124322, and 129262). M. Kivimäki is supported by the Bupa foundation specialist research grant, the UK, and A.S.-M. by a European Young Investigator award from the European Science Foundation.

None.

Disclosures

References

- Scheier MF, Matthews KA, Owens JF, Magovern GJ Sr, Lefebvre RC, Abbott RA, Carver CS. Dispositional optimism and recovery from coronary artery bypass surgery: the beneficial effects on physical and psychological well-being. *J Pers Soc Psychol.* 1989;57:1024–1040.
- Scheier MF, Matthews KA, Owens JF, Schulz R, Bridges MW, Magovern GJ, Carver CS. Optimism and rehospitalization after coronary artery bypass graft surgery. *Arch Intern Med.* 1999;159:829–835.

- Peterson C, Seligman ME, Vaillant GE. Pessimistic explanatory style is a risk factor for physical illness: a thirty-five-year longitudinal study. J Pers Soc Psychol. 1988;55:23–27.
- Giltay EJ, Geleijnse JM, Zitman FG, Hoekstra T, Schouten EG. Dispositional optimism and all-cause and cardiovascular mortality in a prospective cohort of elderly Dutch men and women. *Arch Gen Psychiatry*. 2004;61:1126–1135.
- Tindle HA, Chang YF, Kuller LH, Manson JE, Robinson JG, Rosal MC, Siegle GJ, Matthews KA. Optimism, cynical hostility, and incident coronary heart disease and mortality in the Women's Health Initiative. *Circulation*. 2009;120:656–662.
- Scheier MF, Carver CS, Bridges MW. Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. J Pers Soc Psychol. 1994;67: 1063–1078.
- Kivimaki M, Vahtera J, Elovainio M, Helenius H, Singh-Manoux A, Pentti J. Optimism and pessimism as predictors of change in health after death or onset of severe illness in family. *Health Psychol.* 2005;24: 413–421.
- Kivimaki M, Gimeno D, Ferrie JE, Batty GD, Oksanen T, Jokela M, Virtanen M, Salo P, Akbaraly TN, Elovainio M, Pentti J, Vahtera J. Socioeconomic position, psychosocial work environment and cerebrovascular disease among women: the Finnish public sector study. *Int J Epidemiol.* 2009;38:1265–1271.
- Vahtera J, Kivimaki M, Hublin C, Korkeila K, Suominen S, Paunio T, Koskenvuo M. Liability to anxiety and severe life events as predictors of new-onset sleep disturbances. *Sleep*. 2007;30:1537–1546.
- Giltay EJ, Zitman FG, Kromhout D. Dispositional optimism and the risk of depressive symptoms during 15 years of follow-up: the Zutphen elderly study. J Affect Disord. 2006;91:45–52.
- Larson SL, Owens PL, Ford D, Eaton W. Depressive disorder, dysthymia, and risk of stroke: thirteen-year follow-up from the Baltimore epidemiologic catchment area study. *Stroke*. 2001;32:1979–1983.
- Kubzansky LD, Sparrow D, Vokonas P, Kawachi I. Is the glass half empty or half full? A prospective study of optimism and coronary heart disease in the normative aging study. *Psychosom Med.* 2001;63:910–916.
- Schulz R, Bookwala J, Knapp JE, Scheier M, Williamson GM. Pessimism, age, and cancer mortality. *Psychol Aging*. 1996;11:304–309.
- MacMahon S, Rodgers A. Blood pressure, antihypertensive treatment and stroke risk. J Hypertens Suppl. 1994;12:S5–S14.





Low Pessimism Protects Against Stroke: The Health and Social Support (HeSSup) Prospective Cohort Study

Hermann Nabi, Markku Koskenvuo, Archana Singh-Manoux, Jyrki Korkeila, Sakari Suominen, Katariina Korkeila, Jussi Vahtera and Mika Kivimäki

Stroke. 2010;41:187-190; originally published online November 5, 2009; doi: 10.1161/STROKEAHA.109.565440 Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231 Copyright © 2009 American Heart Association, Inc. All rights reserved. Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at: http://stroke.ahajournals.org/content/41/1/187

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the Permissions and Rights Question and Answer document.

Reprints: Information about reprints can be found online at: http://www.lww.com/reprints

Subscriptions: Information about subscribing to *Stroke* is online at: http://stroke.ahajournals.org//subscriptions/