

Cerebral Aneurysm Morphology Before and After Rupture

Nationwide Case Series of 29 Aneurysms

Torbjørn Øygard Skodvin, medical student; Liv-Hege Johnsen, MD; Øivind Gjertsen, MD; Jørgen Gjernes Isaksen, MD, PhD; Angelika Sorteberg, MD, PhD

Background and Purpose—Using postrupture morphology to predict rupture risk of an intracranial aneurysm may be inaccurate because of possible morphological changes at or around the time of rupture. The present study aims at comparing morphology from angiograms obtained prior to and just after rupture and to evaluate whether postrupture morphology is an adequate surrogate for rupture risk.

Methods—Case series of 29 aneurysms from a nationwide retrospective data collection. Two neuroradiologists who were blinded to pre- versus postrupture images assessed predefined morphological parameters independently and reached consensus regarding all measurements. Prerupture morphology and respective changes after rupture were quantified and linked to risk factors and to the risk of rupture according to the PHASES (population, hypertension, age, size of aneurysm, earlier subarachnoid hemorrhage from another aneurysm, site of aneurysm) and unruptured intracranial aneurysm treatment (UIAT) scores.

Results—All 1-dimensional parameter medians were significantly larger after rupture, except neck diameter. Number of aneurysms with daughter sacs was 9 (31%) before and 17 (59%) after rupture ($P=0.005$). Aneurysm growth from the images prior to and just after rupture increased with the time elapsed between images. Aneurysms in patients with hypertension were significantly larger at diagnosis. Prerupture morphology did not differ in relation to smoke status. Clinical risk factors were not significantly associated with morphological change.

Conclusions—The changes in aneurysm morphology observed after rupture reflect the compound effect of time with successive growth and formation of irregularities and the impact of rupture per se. Postrupture morphology should not be considered an adequate surrogate for the prerupture morphology in the evaluation of rupture risk. (*Stroke*. 2017;48:880-886. DOI: 10.1161/STROKEAHA.116.015288.)

Key Words: cerebral angiography ■ intracranial aneurysm ■ morphology ■ risk factors ■ rupture

Subarachnoid hemorrhage (SAH) causes loss of potential life years at a proportion similar to that of ischemic stroke and intracerebral hemorrhage.^{1,2} The prevalence of intracranial aneurysms (IAs) is estimated to be around 2% to 3.5% in a normal population.^{3,4} Ruptured aneurysms are the source of 80% of SAH. The SAH incidence is 10 per 100 000 person-years, implying that many IAs never rupture.⁵ The optimal management of a diagnosed, unruptured IA remains controversial, and the risks of preventive intervention have to be weighed against the unknown risk of rupture for the individual aneurysm.⁶ Therefore, tools have been developed to establish the risk of IA rupture (such as the PHASES [population, hypertension, age, size of aneurysm, earlier SAH from another aneurysm, site of aneurysm] risk factor score⁷) and to ease the decision-making in the management of unruptured IAs (such as unruptured IA treatment score [UIATS]⁸).

Aneurysm size is often applied in the clinical decision-making because it has been shown to be a significant predictor of

rupture.⁷ On the contrary, rupture is a multifactorial end point and cannot be exactly determined by aneurysm size alone.⁹ Hence, larger aneurysms carry a higher risk of rupture, but, nevertheless, a large amount of SAHs are caused by small aneurysms.¹⁰⁻¹² Aneurysm morphology expressed as aspect ratio and bottleneck factor determines the hemodynamics that may affect the risk of rupture. Several studies showed significant morphological differences between ruptured and unruptured aneurysms.⁹ However, applying results from these studies on unruptured aneurysms to determine their risk of rupture critically relies on the premise that postrupture morphology is not significantly different from that prior to rupture.⁹ Recent case reports and small studies with 1 to 13 aneurysms indicate that this assumption does not hold.¹³⁻¹⁶

The aim of the present study is to describe the changes in morphology and morphological indices occurring between images obtained prior to and just after rupture of an aneurysm and to evaluate whether postrupture morphology is an adequate surrogate for risk of rupture.

Received September 7, 2016; final revision received January 13, 2017; accepted January 20, 2017.

From the UiT The Arctic University of Norway, Tromsø (T.Ø.S., J.G.I.); University Hospital of Northern Norway, Tromsø (L.-H.J., J.G.I.); Oslo University Hospital Rikshospitalet, Norway (Ø.G., A.S.); and Institute of Clinical Medicine, University of Oslo, Norway (A.S.).

The online-only Data Supplement is available with this article at <http://stroke.ahajournals.org/lookup/suppl/doi:10.1161/STROKEAHA.116.015288/-/DC1>.

Correspondence to Torbjørn Øygard Skodvin, Varden 197, 9018 Tromsø, Norway, Europe. E-mail torbjorn.skodvin@gmail.com

© 2017 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.116.015288

Patients and Methods

Study Design, Patient Selection, and Data Extraction

The study is a nationwide retrospective data collection from the 4 neurosurgical centers treating all IA and SAH in Norway. We searched electronic health records using codes from the *International Classification of Diseases*, version 10, to identify patients diagnosed with unruptured IA who later were hospitalized with SAH. We created lists of patients diagnosed with I67.1 (cerebral aneurysm, unruptured) between October 1, 2003, and October 1, 2013. Of these patients, we identified those being subsequently hospitalized with I60.0–I60.9 (nontraumatic SAH).

For the identified patients, we recorded age, sex, date of diagnosis of unruptured IA, date admitted for SAH, and the reason for the aneurysm being conservatively managed. We also retrieved known risk factors, such as hypertension, smoking, connective tissue disease, polycystic kidney disease, family history, and prior aneurysmal SAH. We determined the PHASES⁷ and UIAT⁸ scores in all patients. The latest available prerule and the first available postrupture angiograms were retrieved, from which several quantitative and qualitative features were determined. Patients were excluded if they had previous treatment of the aneurysm of interest, multiple aneurysms of which the ruptured aneurysm was difficult to identify, or image quality precluding reliable aneurysm measurements. Fusiform aneurysms were excluded.

The study was approved by the Northern Norway Regional Committee for Medical Research Ethics, which decided the study to be exempt from patient consent. The study is reported according to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement for observational studies.¹⁷

Measurement Process

Two neuroradiologists residing in different centers assessed the aneurysms using measuring tools in Siemens syngo.via and syngo InSpace (Siemens Healthcare, Erlangen, Germany). The 2 observers measured all aneurysms independently according to a strictly defined measurement protocol, blinded to each other's results and with no prior information about aneurysm rupture state.

Initially, the observers interactively evaluated 3-dimensional volume rendering technique images for general morphology, such as smooth/irregular and numbers of daughter sacs (Figure 1A). The aneurysm neck was identified and multiplanar cursors were aligned to define the aneurysm neck. The aneurysm was rotated until the maximum length and diameters were revealed. The resulting volume rendering technique projection was then converted to a thin-slice maximum-intensity picture on which measurements were performed (Figure 1B).

We calculated intraclass correlation coefficient to assess the absolute interrater agreement.¹⁸ According to conservative criteria, values >0.81 represent substantial reliability.¹⁹ Mean intraclass correlation coefficient for all parameters except 2 was 0.88. Of the 2 with lower intraclass correlation coefficient, 1 parameter (minimal size) was excluded from further analyses, whereas the other (Neck) was redefined to increase precision. The final measurement guide is described below (see Parameter Definitions).

The mean values between observers were chosen when interrater difference was <2 mm. In cases of ≥2 mm differences and for the redefined Neck parameter, values were settled by consensus.

Parameter Definitions

Size Measurements

Definitions are illustrated in Figure 1C and 1D. All parameters must be fitted within the aneurysm sac. Maximal size is the maximal distance between any 2 points in the aneurysm sac, including the neck plane. Neck size is the largest observed diameter of the neck plane. Height is the orthogonal distance between the neck plane center and the aneurysm dome. Length is the greatest distance between the neck plane center and any point on the aneurysm dome, not necessarily orthogonal

to the neck plane. Width *L* is the largest diameter that is orthogonal to length. Width *H* is the largest diameter that is orthogonal to height. When comparing digital subtraction angiography with another image modality, the available digital subtraction angiography projections dictated which projections were used from the other modality.

Indices

Aspect ratio was calculated as the ratio between height and neck diameter, and the bottleneck factor was calculated as the ratio between width *L* and neck diameter.²⁰

Volume Approximation

Three of the above measured diameters were used to mathematically approximate the aneurysm volume. The conventional volume formula is $V=4/3 \times \pi(A/2) \times (B/2) \times (C/2)$. We replaced *A*, *B*, and *C* with maximal size, length, and width *L*, respectively. This approximation technique has been shown to underestimate but still correlate with other methods of volume measurements.²¹

Additional Registrations

We registered aneurysm location, relation to parent artery (bifurcation or sidewall aneurysm), surface quality (smooth or irregular), and the presence and number of daughter sacs protruding from the aneurysm wall.

Statistical Analysis

The data were analyzed with Stata for Mac (version 14; StataCorp LP, TX) and SPSS for Windows (version 24; IBM, NY). The variables' distribution was investigated visually with Q-Q plots and numerically with Shapiro–Wilk test and found to be nonparametric. Thus, paired continuous variables were compared using Wilcoxon signed-rank test, and paired nominal variables were compared using McNemar's test. Independent continuous variables were compared using Mann–Whitney *U* test or Kruskal–Wallis test in cases of >2 groups. Categorical variables were compared using Chi-squared test. A *P* value of <0.05 was assumed statistically significant.

Results

The search identified 52 patients with confirmed aneurysmal SAH, originating from aneurysms that were recognized prior to rupture but not repaired. Of these, 23 were excluded (9 were fusiform and 14 because of missing or poor images). The remaining 29 patients were included in the study.

Eight of the 29 patients were men (28%). Mean age at time of SAH was 67 years (SD, 9.3). Median time span between imaging prior to and just after rupture was 12 months (range, 0.3–96 months). The combination of image modalities before and after rupture was computed tomography angiography (CTA) and CTA for 16 patients (56%), CTA and magnetic resonance angiography for 8 patients (28%), CTA and 2D digital subtraction angiography for 2 patients (7%), and magnetic resonance angiography and 2D digital subtraction angiography for 3 patients (10%).

Table 1 summarizes patient and aneurysm characteristics at baseline, as well as risk of rupture expressed by PHASES risk score and treatment recommendations according to the UIAT score. The 5-year risk of rupture exceeded 1% (PHASES ≥5) in 79% of the patients and 5% (PHASES ≥10) in 38% of the patients. The UIAT score was indeterminate in 35% of the cases and favored conservative management in 31% of the patients. The UIAT score favoring aneurysm repair tended to be higher for aneurysms that ruptured within 3 months (median, 14; range, 5–16) compared with aneurysms with longer time




Figure 1. Aneurysm measurement method. **A**, Volume rendering technique (VRT) image for assessment of general morphology and identification of optimal measurement planes. **B**, Maximum intensity picture (MIP) for measurements. **C** and **D**, Illustrations of parameter definitions.

between images (median, 10; range 5–20; $P=0.07$). Table in the [online-only Data Supplement](#) provides aneurysm location, maximal aneurysm size, PHASES and UIAT scores, and the reason for not performing aneurysm repair for each individual aneurysm.

Morphological Change

Table 2 lists the morphological parameters from imaging prior to and just after rupture. All 1-dimensional parameter medians were statistically significantly larger after rupture, except neck diameter. Median aspect ratio before rupture was 1.5 (range, 0.8–4.0) compared with 1.9 (range, 0.8–6.7) after rupture ($P=0.008$). Median bottleneck factor was 1.5 (range, 0.9–4.0) before and 1.5 (range, 0.7–6.2) after rupture ($P=0.068$). Number of aneurysms with ≥ 1 daughter sac was 9 (31%) before and 17 (59%) after rupture ($P=0.005$). Figure 2 illustrates a typical change from pre- to postrupture image.

The magnitude of change was clearly dependent on the time elapsed between the image prior to and just after rupture, visualized in Figure 3. Seven aneurysms (24%) ruptured within 3 months after the last image. Though median change in aspect ratio for this subset was only 0.10, the range from -0.45 to 0.95 (corresponding to -18% to 98%) demonstrates that we also observed relatively large morphological change within short time spans. We observed a new daughter sac in 1 of these 7 aneurysms after rupture (number of days between images for this particular aneurysm was 18). Change in the

morphological indices, however, was not dependent on time elapsed between images.

Morphology and Change in Relation to Clinical Risk Factors

In hypertensive patients, aneurysms were significantly larger at diagnosis (median maximum size, 12.85 mm [range, 3.4–12.3] in hypertensive versus 6.95 mm [2.8–33.4] in nonhypertensive; $P=0.041$). Indices and presence of daughter sacs were not significantly different. Changes in morphology from before to after rupture were not significant, except for neck diameter, which tended to increase in hypertensive patients (0.3 mm [-1.8 to 3.7]) and decrease (-0.4 mm [-2.9 to 0.5]) in nonhypertensive patients ($P=0.047$).

Between current and former/never smokers, there was neither a significant difference in morphology prior to rupture nor a significant change in morphology after rupture. These findings were also true when excluding aneurysms that ruptured after a short observation period (<3 months).

Discussion

The core finding of the present study is that aneurysm morphology had changed between imaging prior to and just after aneurysm rupture. The observed changes increased with the time elapsed between imaging, though gross changes also occurred within short time spans. Change occurred in a non-uniform manner, signified by changes in aspect ratio and fraction of aneurysms with daughter sacs.

Table 1. Patient and Aneurysm Characteristics

Characteristic	No. (%)
Patients	
No. of patients	29 (100)
Sex, male	8 (28)
Age at time of SAH, mean (\pm SD), y	67.2 (9.3)
Hypertension	21 (72)
Multiple aneurysms	12 (41)
Smoking	
Current	13 (45)
Former	3 (10)
Never	13 (45)
PHASES 5-year rupture risk, median (range)	
>1% risk, n (%)	23 (79)
>5% risk, n (%)	11 (38)
UIAT score	
Favored repair	10 (35)
Indeterminate	9 (31)
Favored conservative management	10 (35)
Aneurysms	
No. of aneurysms	29 (100)
Location*	
Anterior	26
Posterior	3
Time between images, median (range), mo	12 (0.26–96.2)

PHASES indicates population, hypertension, age, size of aneurysm, earlier SAH from another aneurysm, site of aneurysm; SAH, subarachnoid hemorrhage; and UIAT, unruptured intracranial aneurysm treatment.

*Anterior: anterior cerebral artery, anterior communicating artery, middle cerebral artery and internal carotid artery. Posterior: basilar artery, posterior cerebral artery, posterior communicating artery.

Postrupture Morphology as Surrogate for Prerupture Morphology

The present findings do not support the assumption that postrupture morphology is representative for the prerupture morphology.⁹ The fraction of aneurysms with blebs increased from 31% before to 59% after rupture. Consistent with the present study, a recent literature review reported that 17 of 23 aneurysms increased in size around time of rupture, and a case series in the same work showed presence of new daughter sacs after rupture in 5 out of 6 patients.¹⁶ However, aneurysm morphology just after rupture will be subjected to the impact of the rupture per se plus any change that may have occurred along the evolution of the specific aneurysm or even in the short time span between rupture and postrupture imaging. In a meta-analysis including 4972 unruptured aneurysms, 9% of aneurysms enlarged within a mean follow-up time of 2.8 patient-years.²² Accordingly, our data do not reveal what occurs during the exact moment of rupture, but rather support the notion that aneurysms grow over time, with periods with and without growth, and an inconstant risk of rupture over time.^{23–25}

Table 2. Morphological Parameters Before and After Rupture

	Before Rupture, Median (Range)	After Rupture, Median (Range)	P Values*
1D parameters, mm			
Maximal diameter	10.0 (2.8–33.4)	12.1 (3.5–40.2)	<0.001†
Neck diameter	5.6 (1.9–12.8)	5.4 (2.1–13.1)	0.79
Length	9.6 (2.7–25.4)	11.1 (3.5–40.2)	0.003†
Width <i>L</i>	8.6 (2.5–28.0)	9.4 (2.1–37.0)	0.024†
Height	9.6 (2.7–25.4)	9.1 (2.6–40.2)	0.035†
Width <i>H</i>	8.4 (2.5–28.0)	9.6 (2.1–37.0)	0.002†
2D parameters			
Aspect ratio	1.5 (0.8–4.0)	1.9 (0.8–6.7)	0.008†
Bottleneck factor	1.5 (0.9–4.0)	1.5 (0.7–6.2)	0.069
3D parameters			
Approximated volume, cm ³	0.50 (0.01–11.2)	0.57 (0.02–31.3)	0.001†
Wall characteristics			
Irregular, n (%)	10 (35)	13 (45)	0.25‡
No. of blebs, median (range)	0 (0–2)	1 (0–5)	0.001†
Aneurysms with blebs, n (%)	9 (31)	17 (59)	0.005†

*P value for the difference before and after rupture; Wilcoxon signed-rank test.

†P values <0.05 considered statistically significant.

‡Related samples McNemar test.

The low rate of rupture of small aneurysms in the International Study of Intracranial Aneurysms led some authors to speculate that aneurysms shrink after rupture.^{14,26} Aneurysms in the present study that ruptured within 3 months after the prerupture image showed less change in morphology (or even shrunk in some of the parameters) than those that had ruptured after longer time intervals. One could assume that the changes in this subgroup were more subjected to the effect of the rupture per se than those we observed in the other aneurysms. Though the changes are too small to rule out measurement uncertainty, one can speculate that rupture may cause a slight deflation of aneurysms. Three of the aneurysms ruptured after 9, 16, and 22 months after the last prerupture scan, respectively, and also showed a decrease in maximum size; one could speculate that these were stable aneurysms without growth during the time span and that the observed change was caused by the rupture. However, the number of aneurysms is too small to allow for conclusions.

Risk of Rupture

An aspect ratio >1.6 has been considered as a predictor for future rupture and is also included in the UIAT score.⁸ We are not aware of suggested cutoff values for bottleneck factor. Although such thresholds have been criticized and are affected by measurement methodology,²⁰ we note with interest that 15 (52%) of the 29 aneurysms in our material fell below these limits before rupture, whereas 10 (35%) did so after rupture. Because the mean values increased for all parameters except




Figure 2. Volume rendering technique (VRT) images of 1 aneurysm before and after rupture. Height and maximal size measurements are shown. **A**, Before rupture. **B**, After rupture. Maximal size is increased and a daughter sac has developed.

neck diameter, any change would tend to increase aspect ratio and bottleneck factor. Thus, the higher aspect ratio and bottleneck factor seen in ruptured aneurysms in other studies may simply be the effect of change over time or the rupture itself.²⁷

Maximal aneurysm diameter is perhaps the most common denominator for determining the risk of rupture and is incorporated into the PHASES⁷ and UIAT⁸ scores. The fact that even small aneurysms rupture is well established.^{28,29} Close

to one third (28%) of our aneurysms were <7 mm and still ruptured. The size of our aneurysms did not predict the time span to rupture. One reason may be an overrepresentation of small aneurysms within the population of unruptured IAs and another may be that aneurysms grow in a nonlinear fashion.^{23,24}

Size is only 1 indicator of the multifactorial causes for aneurysm rupture. PHASES and UIAT scores incorporate a wide array of factors supposedly influencing the risk of




Figure 3. Difference between pre- and postrupture measurements of the 1-dimensional parameters maximal size, height, neck, and length, categorized in accordance with the time elapsed between images. *P* values from independent samples Kruskal–Wallis test.

aneurysm rupture; still, in merely about one third of our cases, the UIAT score would have favored aneurysm repair. In 20% of cases, the PHASES score indicated a 5-year cumulative rupture risk of <1%. On the contrary, the UIAT score favored repair or suggested special consideration because of indetermination regarding treatment in 70% of patients. According to PHASES, 38% of the patients exceeded a 5-year risk of rupture of 5%. Though our retrospective study aims to compare pre- and postrupture imaging, we note that the majority of the included patients may have required treatment.

Aneurysm growth is a strong risk factor for rupture,^{22,24,30–32} possibly increasing risk 12- to 24-fold.^{30,31} Growth rate and risk of growth increases with increased aneurysm size. However, growth can occur at all aneurysm sizes, warranting follow-up imaging of conservatively managed aneurysms, including aneurysms <7 mm.^{24,25,27,29–31} In a systematic review of 30 unruptured aneurysms <7 mm followed with serial imaging for a median of 6.5 years, 27 (90%) enlarged before rupture.²⁴ Thus, assuming that at least substantial parts of the changes we observe in our study are prerupture changes, our study sample consists of aneurysms of a high rupture risk. Still, aneurysm growth is only one marker of increased risk, and rupture can occur without growth.^{24,25}

Smoking and hypertension are other well-established independent risk factors for aneurysm rupture.^{8,33} Morphological changes, however, were similar for patients with and without these risk factors. Thus, in our material, the presence or absence of smoking and hypertension did not influence whether postrupture morphology was representative of that prior to rupture.

Studies comparing unruptured aneurysms with aneurysms presenting after rupture have generated important hypotheses about pathophysiology and risk factors for growth and rupture.³⁴ With the addition of the present study to existing data, we argue that the postrupture morphology should not be considered a good surrogate in the evaluation of risk of rupture. Morphological and hemodynamic rupture predictors should be validated in studies of prerupture aneurysms.

Limitations

The present material is subjected to selection bias because the included patients were selected to conservative management, except those who either refused treatment or experienced SAH while waiting for aneurysm repair. Several factors can affect the rupture risk: the included patients are somewhat older and possibly more comorbid than the expected average of SAH patients. The fraction of smokers in our material is somewhat lower than the country's average (less than half versus 2 thirds, respectively), and the fraction of patients with multiple aneurysms is higher than what is commonly found in clinical series.

A length time bias may pertain to the included aneurysms because other more rupture-prone aneurysms might have ruptured early on in their pathogenesis, never being diagnosed before rupture. Also, patients with a recognized IA that succumbed to their aneurysm without reaching a hospital are not part of this study. Still, we regard the external validity as high because none of the clinical risk factors were statistically significantly associated with morphological change. However,

the retrospective nature of the study reduces the accuracy of the patient risk factors recorded. The study sample is small and does not allow for definite conclusions, but pre- and postrupture angiograms of IAs are exceedingly rare, making adequately powered enquiries into this matter difficult. This study, thus, contributes to shed light on an area that is rarely available for investigation.

The neck diameter definition used in this study is the maximal neck size. This definition provided the highest interrater reliability, but reduces comparability with studies using average or minimal neck diameter definitions. To answer our study question, the interrater reliability was paramount. Manual measurements can introduce interrater discrepancies. Strict parameter definitions guided the measuring process to counter error. The intraclass correlation coefficient demonstrated substantial agreement between the 2 raters.

Finally, 45% of the cases were evaluated with different image modalities, introducing technical limitations in measurement precision. However, other studies have shown that different modalities can be reliably compared.³⁵ Moreover, image quality, CTA and magnetic resonance angiography slice thickness, as well as radiocontrast filling effects introduce variability. This variability may be assumed to be of the same magnitude like the one meeting the clinician in every day practice when evaluating serial imaging in a patient. The observed changes in the present study are of such a magnitude that they still would be consistent even after considering a margin for measurement errors.

Conclusions

Aneurysm morphology was significantly different after rupture as compared with that before rupture. To an extent, changes had occurred in a nonuniform manner. The changes observed after rupture reflect the compound effect of time with successive growth and formation of irregularities and the impact of rupture per se. Postrupture morphology should not be considered a good surrogate in the evaluation of risk of rupture.

Acknowledgments

We thank Rune Grov Eilertsen (University Hospital of Northern Norway, Tromsø, Norway), Christian A. Helland (Haukeland University Hospital, Bergen, Norway), Ole Solheim (St. Olav Hospital, Trondheim, Norway), and Wilhelm Sorteberg (Oslo University Hospital Rikshospitalet, Norway) for participating in the data collection.

Sources of Funding

UiT The Arctic University of Norway funded the study. The funder had no role in the study design, data collection, data analysis, data interpretation, writing of the article or decision to submit for publication.

Disclosures

None.

References

1. Johnston SC, Selvin S, Gress DR. The burden, trends, and demographics of mortality from subarachnoid hemorrhage. *Neurology*. 1998;50:1413–1418.

2. Nieuwkamp DJ, Setz LE, Algra A, Linn FH, de Rooij NK, Rinkel GJ. Changes in case fatality of aneurysmal subarachnoid haemorrhage over time, according to age, sex, and region: a meta-analysis. *Lancet Neurol*. 2009;8:635–642. doi: 10.1016/S1474-4422(09)70126-7.
3. Müller TB, Sandvei MS, Kvistad KA, Rydland J, Håberg A, Vik A, et al. Unruptured intracranial aneurysms in the Norwegian Nord-Trøndelag Health Study (HUNT): risk of rupture calculated from data in a population-based cohort study. *Neurosurgery*. 2013;73:256–261, discussion 260, quiz 261. doi: 10.1227/01.neu.0000430295.23799.16.
4. Vlak MH, Algra A, Brandenburg R, Rinkel GJ. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *Lancet Neurol*. 2011;10:626–636. doi: 10.1016/S1474-4422(11)70109-0.
5. Lindekleiv HM, Njølstad I, Ingebrigtsen T, Mathiesen EB. Incidence of aneurysmal subarachnoid hemorrhage in Norway, 1999–2007. *Acta Neurol Scand*. 2011;123:34–40. doi: 10.1111/j.1600-0404.2010.01336.x.
6. Brown RD Jr, Broderick JP. Unruptured intracranial aneurysms: epidemiology, natural history, management options, and familial screening. *Lancet Neurol*. 2014;13:393–404. doi: 10.1016/S1474-4422(14)70015-8.
7. Greving JP, Wermer MJ, Brown RD Jr, Morita A, Juvela S, Yonekura M, et al. Development of the PHASES score for prediction of risk of rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *Lancet Neurol*. 2014;13:59–66. doi: 10.1016/S1474-4422(13)70263-1.
8. Etminan N, Brown RD Jr, Beseoglu K, Juvela S, Raymond J, Morita A, et al. The unruptured intracranial aneurysm treatment score: a multidisciplinary consensus. *Neurology*. 2015;85:881–889. doi: 10.1212/WNL.0000000000001891.
9. Zanaty M, Chalouhi N, Tjoumakaris SI, Fernando Gonzalez L, Rosenwasser RH, Jabbour PM. Aneurysm geometry in predicting the risk of rupture. A review of the literature. *Neurol Res*. 2014;36:308–313. doi: 10.1179/1743132814Y.00000000327.
10. Weir B. Unruptured intracranial aneurysms: a review. *J Neurosurg*. 2002;96:3–42. doi: 10.3171/jns.2002.96.1.0003.
11. Joo SW, Lee SI, Noh SJ, Jeong YG, Kim MS, Jeong YT. What is the significance of a large number of ruptured aneurysms smaller than 7 mm in diameter? *J Korean Neurosurg Soc*. 2009;45:85–89. doi: 10.3340/jkns.2009.45.2.85.
12. Forget TR Jr, Benitez R, Veznedaroglu E, Sharan A, Mitchell W, Silva M, et al. A review of size and location of ruptured intracranial aneurysms. *Neurosurgery*. 2001;49:1322–1325, discussion 1325.
13. Kono K, Tomura N, Yoshimura R, Terada T. Changes in wall shear stress magnitude after aneurysm rupture. *Acta Neurochir (Wien)*. 2013;155:1559–1563. doi: 10.1007/s00701-013-1773-2.
14. Rahman M, Ogilvy CS, Zipfel GJ, Derdeyn CP, Siddiqui AH, Bulsara KR, et al. Unruptured cerebral aneurysms do not shrink when they rupture: multicenter collaborative aneurysm study group. *Neurosurgery*. 2011;68:155–160, discussion 160. doi: 10.1227/NEU.0b013e3181ff357c.
15. Schneiders JJ, Marquering HA, van den Berg R, VanBavel E, Velthuis B, Rinkel GJ, et al. Rupture-associated changes of cerebral aneurysm geometry: high-resolution 3D imaging before and after rupture. *AJNR Am J Neuroradiol*. 2014;35:1358–1362. doi: 10.3174/ajnr.A3866.
16. Yi J, Zielinski D, Chen M. Cerebral aneurysm size before and after rupture: case series and literature review. *J Stroke Cerebrovasc Dis*. 2016;25:1244–1248. doi: 10.1016/j.jstrokecerebrovasdis.2016.01.031.
17. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med*. 2007;147:573–577.
18. Gisev N, Bell JS, Chen TF. Interrater agreement and interrater reliability: key concepts, approaches, and applications. *Res Social Adm Pharm*. 2013;9:330–338. doi: 10.1016/j.sapharm.2012.04.004.
19. Shrout PE. Measurement reliability and agreement in psychiatry. *Stat Methods Med Res*. 1998;7:301–317.
20. Lauric A, Baharoglu MI, Malek AM. Ruptured status discrimination performance of aspect ratio, height/width, and bottleneck factor is highly dependent on aneurysm sizing methodology. *Neurosurgery*. 2012;71:38–45. doi: 10.1227/NEU.0b013e3182503bf9.
21. Sadato A, Hayakawa M, Tanaka T, Hirose Y. Comparison of cerebral aneurysm volumes as determined by digitally measured 3D rotational angiography and approximation from three diameters. *Interv Neuroradiol*. 2011;17:154–158.
22. Backes D, Rinkel GJ, Laban KG, Algra A, Vergouwen MD. Patient- and aneurysm-specific risk factors for intracranial aneurysm growth: a systematic review and meta-analysis. *Stroke*. 2016;47:951–957. doi: 10.1161/STROKEAHA.115.012162.
23. Koffijberg H, Buskens E, Algra A, Wermer MJ, Rinkel GJ. Growth rates of intracranial aneurysms: exploring constancy. *J Neurosurg*. 2008;109:176–185. doi: 10.3171/JNS/2008/109/8/0176.
24. Chmaysani M, Rebeiz JG, Rebeiz TJ, Batjer HH, Bendok BR. Relationship of growth to aneurysm rupture in asymptomatic aneurysms ≤ 7 mm: a systematic analysis of the literature. *Neurosurgery*. 2011;68:1164–1171, discussion 1171. doi: 10.1227/NEU.0b013e31820edbd3.
25. Bor AS, Tiel Groenestege AT, terBrugge KG, Agid R, Velthuis BK, Rinkel GJ, et al. Clinical, radiological, and flow-related risk factors for growth of untreated, unruptured intracranial aneurysms. *Stroke*. 2015;46:42–48. doi: 10.1161/STROKEAHA.114.005963.
26. Sonobe M, Yamazaki T, Yonekura M, Kikuchi H. Small unruptured intracranial aneurysm verification study: SUAVE study, Japan. *Stroke*. 2010;41:1969–1977. doi: 10.1161/STROKEAHA.110.585059.
27. Żyłkowski J, Kunert P, Jaworski M, Rosiak G, Marchel A, Rowiński O. Changes of size and shape of small, unruptured intracranial aneurysms in repeated computed tomography angiography studies. *Wideochir Inne Tech Maloinwazyjne*. 2015;10:178–188. doi: 10.5114/wiitm.2015.52707.
28. Froelich JJ, Neilson S, Peters-Wilke J, Dubey A, Thani N, Erasmus A, et al. Size and location of ruptured intracranial aneurysms: a 5-year clinical survey. *World Neurosurg*. 2016;91:260–265. doi: 10.1016/j.wneu.2016.04.044.
29. Serrone JC, Tackla RD, Gozal YM, Hanseman DJ, Gogela SL, Vuong SM, et al. Aneurysm growth and de novo aneurysms during aneurysm surveillance. *J Neurosurg*. 2016;125:1374–1382. doi: 10.3171/2015.12.JNS151552.
30. Villablanca JP, Duckwiler GR, Jahan R, Tateshima S, Martin NA, Frazee J, et al. Natural history of asymptomatic unruptured cerebral aneurysms evaluated at CT angiography: growth and rupture incidence and correlation with epidemiologic risk factors. *Radiology*. 2013;269:258–265. doi: 10.1148/radiol.13121188.
31. Teo M, St George EJ. Radiologic surveillance of untreated unruptured intracranial aneurysms: a single surgeon's experience. *World Neurosurg*. 2016;90:20–28. doi: 10.1016/j.wneu.2016.02.008.
32. Brinjikji W, Zhu YQ, Lanzino G, Cloft HJ, Murad MH, Wang Z, et al. Risk factors for growth of intracranial aneurysms: a systematic review and meta-analysis. *AJNR Am J Neuroradiol*. 2016;37:615–620. doi: 10.3174/ajnr.A4575.
33. Sandvei MS, Romundstad PR, Müller TB, Vatten L, Vik A. Risk factors for aneurysmal subarachnoid hemorrhage in a prospective population study: the HUNT study in Norway. *Stroke*. 2009;40:1958–1962. doi: 10.1161/STROKEAHA.108.539544.
34. Xiang J, Natarajan SK, Tremmel M, Ma D, Mocco J, Hopkins LN, et al. Hemodynamic-morphologic discriminants for intracranial aneurysm rupture. *Stroke*. 2011;42:144–152. doi: 10.1161/STROKEAHA.110.592923.
35. Goubergrits L, Schaller J, Kertzscher U, Petz Ch, Hege HC, Spuler A. Reproducibility of image-based analysis of cerebral aneurysm geometry and hemodynamics: an in-vitro study of magnetic resonance imaging, computed tomography, and three-dimensional rotational angiography. *J Neurol Surg A Cent Eur Neurosurg*. 2013;74:294–302. doi: 10.1055/s-0033-1342937.

Cerebral Aneurysm Morphology Before and After Rupture: Nationwide Case Series of 29 Aneurysms

Torbjørn Øygaard Skodvin, Liv-Hege Johnsen, Øivind Gjertsen, Jørgen Gjernes Isaksen and Angelika Sorteberg

Stroke. 2017;48:880-886; originally published online March 6, 2017;
doi: 10.1161/STROKEAHA.116.015288

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2017 American Heart Association, Inc. All rights reserved.
Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/48/4/880>

An erratum has been published regarding this article. Please see the attached page for:
</content/48/4/e121.full.pdf>

Data Supplement (unedited) at:

<http://stroke.ahajournals.org/content/suppl/2017/03/06/STROKEAHA.116.015288.DC1>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Stroke* is online at:
<http://stroke.ahajournals.org/subscriptions/>

Correction to: Cerebral Aneurysm Morphology Before and After Rupture: Nationwide Case Series of 29 Aneurysms

In the article by Skodvin et al, “Cerebral Aneurysm Morphology Before and After Rupture: Nationwide Case Series of 29 Aneurysms,” which published online ahead of print March 6, 2017, and appeared in the April 2017 issue of the journal (*Stroke*. 2017;48:880–886. DOI: 10.1161/STROKEAHA.116.015288), a correction is needed.

On page 883, in Table 2 footnotes, “†*P* values <0.05 considered statistically significant and given in bold,” has been changed to read “†*P* values <0.05 considered statistically significant.”

This correction has been made to the current online version of the article, which is available at <http://stroke.ahajournals.org/content/48/4/e121>.

Online Supplement

CEREBRAL ANEURYSM MORPHOLOGY BEFORE AND AFTER RUPTURE:
NATION-WIDE CASE SERIES OF 29 ANEURYSMS.

Table I. Location, Time Frame, Maximum Size, Risk Factor Scores and Reasons For Not Treating the Individual Aneurysm
References

1
2

Table I. Location, Time Frame, Maximum Size, Risk Factor Scores and Reasons For Not Treating the Individual Aneurysm.

Location	Time between last imaging and rupture (months)	Maximum size when diagnosed (mm)	PHASES score	UIATS*			Reason for not treating
				Favors repair	Favors conservative management	Difference	
Middle cerebral artery	12.0	6.2	3	5	13	-8	High co-morbidity
	22.3	12.3	5	6	11	-4	Patient refused treatment
	24.6	7.0	6	9	9	0	Patient considered too old (70 years)
	16.5	7.2	6	7	10	-3	Aneurysm deemed too small
	12.6	27.4	13	9	16	-7	Considered ineligible for treatment
	79.1	6.6	4	5	10	-5	Patient refused treatment
	8.9	33.4	13	13	16	-3	Considered ineligible for treatment
	1.5	12.2	9	13	9	4	Bled before decision was made
	7.7	3.4	3	6	8	-2	Patient considered too old (70 years)
	52.5	7.8	5	13	7	4	Patient lost to follow-up
23.9	2.8	3	11	6	5	Aneurysm deemed too small	
Pericallosal artery	1.5	4.7	4	5	6	-1	Aneurysm overlooked at initial scan
Anterior communicating artery	17.6	9.3	8	11	10	1	Aneurysm deemed too small
	14.0	30.3	16	17	17	0	Considered ineligible for treatment
	1.6	9.9	7	16	7	9	Bled while waiting for treatment
	0.3	12.9	12	14	11	3	Bled while waiting for treatment
	6.5	3.3	5	10	6	4	Aneurysm deemed too small
	96.2	8.4	8	12	9	3	Patient considered too old (69 years)
52.3	17.3	11	15	14	1	High co-morbidity	
Posterior communicating artery	37.2	6.7	6	5	10	-5	Considered ineligible for treatment
	57.2	10.0	11	8	12	-4	Considered ineligible for treatment
	9.7	4.5	4	9	8	1	Patient refused treatment
	0.6	13.0	12	14	12	2	Bled while waiting for treatment
	10.3	14.2	12	12	12	0	Patient refused treatment
Basilar tip	1.8	14.3	12	15	12	3	Bled while waiting for treatment
	11.6	11.1	8	13	15	-2	Considered ineligible for treatment
	0.3	14.8	11	13	9	4	Bled while waiting for treatment
Internal carotid artery	21.2	30.1	11	13	16	-3	Considered ineligible for treatment
	9.0	15.7	8	20	9	11	Bled while waiting for treatment

PHASES indicates 5-year absolute risk of aneurysm of rupture¹; UIATS, The unruptured intracranial aneurysm treatment score.²

* UIATS difference +- 2 is considered “not definite”, and either management approach could be supported.²

References

1. Greving JP, Wermer MJ, Brown RD, Jr., Morita A, Juvela S, Yonekura M, et al. Development of the PHASES score for prediction of risk of rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *Lancet Neurol.* 2014;13:59-66
2. Etminan N, Brown RD, Jr., Beseoglu K, Juvela S, Raymond J, Morita A, et al. The unruptured intracranial aneurysm treatment score: a multidisciplinary consensus. *Neurology.* 2015;85:881-889