

Palliative Care for Hospitalized Patients With Stroke

Results From the 2010 to 2012 National Inpatient Sample

Tarvinder Singh, MD; Steven R. Peters, MD; David L. Tirschwell, MD; Claire J. Creutzfeldt, MD

Background and Purpose—Substantial variability exists in the use of life-prolonging treatments for patients with stroke, especially near the end of life. This study explores patterns of palliative care utilization and death in hospitalized patients with stroke across the United States.

Methods—Using the 2010 to 2012 nationwide inpatient sample databases, we included all patients discharged with stroke identified by *International Classification of Diseases-Ninth Revision* codes. Strokes were subclassified as ischemic, intracerebral, and subarachnoid hemorrhage. We compared demographics, comorbidities, procedures, and outcomes between patients with and without a palliative care encounter (PCE) as defined by the *International Classification of Diseases-Ninth Revision* code V66.7. Pearson χ^2 test was used for categorical variables. Multivariate logistic regression was used to account for hospital, regional, payer, and medical severity factors to predict PCE use and death.

Results—Among 395 411 patients with stroke, PCE was used in 6.2% with an increasing trend over time ($P < 0.05$). We found a wide range in PCE use with higher rates in patients with older age, hemorrhagic stroke types, women, and white race (all $P < 0.001$). Smaller and for-profit hospitals saw lower rates. Overall, 9.2% of hospitalized patients with stroke died, and PCE was significantly associated with death. Length of stay in decedents was shorter for patients who received PCE.

Conclusions—Palliative care use is increasing nationally for patients with stroke, especially in larger hospitals. Persistent disparities in PCE use and mortality exist in regards to age, sex, race, region, and hospital characteristics. Given the variations in PCE use, especially at the end of life, the use of mortality rates as a hospital quality measure is questioned. (*Stroke*. 2017;48:2534-2540. DOI: 10.1161/STROKEAHA.117.016893.)

Key Words: end-of-life ■ inpatients ■ palliative care ■ stroke ■ subarachnoid hemorrhage ■ United States

Stroke is the leading cause of serious long-term disability in the United States and the fifth leading cause of death.^{1,2} The stroke illness trajectory presents a unique challenge that is distinct from most other serious illnesses;³ the presentation is sudden and unexpected, and whereas a small proportion of patients will receive potentially curative acute treatment, patients with severe stroke typically face treatment decisions that leave little time for deliberation and can lead either to an early death in the setting of withdrawal or withholding of life-sustaining interventions or enable survival with a wide range of disability.³ In some cases, survival may be worse than death.⁴ Given their neurological impairment, conversations about goals of care usually occur between providers and surrogate decision makers, rather than with the patient themselves.

These observations highlight the importance of integrating palliative care into the acute stroke care setting.⁵ Palliative care is a multidisciplinary approach to medical care that focuses on improving communication, decision making, and quality of life for patients with serious illness and their families. Early integration of palliative care into acute stroke care has been recently endorsed by the American Heart/American Stroke

Association⁷ and Neurocritical Care Society,⁸ but data remain limited about how to best implement these recommendations and how to measure their benefit.⁹⁻¹¹ Palliative care services are available in an increasing number of US hospitals, but substantial variations remain in access and use across regional, socioeconomic, racial, and ethnic groups.^{12,13}

In an effort to evaluate hospital quality of care, the Center for Medicare and Medicaid Services (CMS) currently uses 30-day mortality after ischemic stroke as a quality of care surrogate¹⁴ and provides access to adjusted rates on the CMS hospital compare website (<https://www.medicare.gov/hospitalcompare/search.html>). Inpatient mortality after stroke varies widely depending on several patient and hospital characteristics, including the hospital's use of do not resuscitate (DNR) orders.^{7,15,16} Use of palliative care after stroke is likely also associated with higher in-hospital and 30-day mortality rates and may not indicate lower hospital quality of care: this is not accounted for in the CMS stroke mortality quality measure.

The overall goal of this study was to characterize current practices around the use of palliative care in a nationally

Received March 2, 2017; final revision received July 14, 2017; accepted July 14, 2017.

From the Department of Neurology, Harborview Medical Center, University of Washington, Seattle.

The online-only Data Supplement is available with this article at <http://stroke.ahajournals.org/lookup/suppl/doi:10.1161/STROKEAHA.117.016893/-/DC1>.

Correspondence to Tarvinder Singh, MD, Department of Neurology, Harborview Medical Center, 325 9th Ave, Seattle, WA 98104. E-mail nanites@uw.edu
© 2017 The Author(s). *Stroke* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the Creative Commons Attribution License (CC-BY), which permits use, distribution, and reproduction in any medium, provided the original work is properly cited.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.117.016893

representative sample of patients with stroke by (1) identifying patient and hospital characteristics associated with palliative care utilization, and (2) assessing how the use of palliative care influences inpatient mortality.

Methods

Database

We performed a retrospective observational study in patients with stroke admitted to US acute care hospitals using discharge data from the publicly available national inpatient sample (NIS), healthcare cost and utilization project, and agency for healthcare research and quality.^{17–19}

The NIS is a cross-sectional, all-payer, inpatient care data set in the United States, consolidated on an annual basis. It is the largest inpatient health data set in the United States. Unweighted, it contains data from >7 million hospital stays from >1000 hospitals each year, which represent a stratified sample of 20% of all nonfederal hospitals. Weighted, it estimates >35 million hospitalizations nationally. Discharge data include demographics, socioeconomic, primary and secondary diagnoses, procedures, and length of stay (LOS). The NIS database contains deidentified information and is exempt from institution review board approval at our institution.

Stroke Data Selection

We identified adult (age, >18 years) stroke admissions from 2010 to 2012 using *International Classification of Diseases-Ninth Revision* (ICD-9) diagnosis codes. Codes 433.X1—occlusion and stenosis of cerebral artery with infarction, 434.X1—occlusion of cerebral artery with infarction, and 436—acute but ill-defined cerebrovascular disease, irrespective of their diagnosis position, were used to identify ischemic strokes. Code 430 (first diagnosis only) was used to identify subarachnoid hemorrhage (SAH) and 431 (first diagnosis only) for intracerebral hemorrhage (ICH). Cases were excluded if there was a concomitant ICD-9 code for traumatic brain injury or rehabilitation stay.²⁰

Demographic and socioeconomic factors were identified from the primary data set. Race/ethnicity had a high degree of missing data compared with other variables because of state suppression or partial reporting by hospitals. We identified individuals with intubation and PEG (percutaneous endoscopic gastrostomy) tube placement separately as proxy for life-prolonging care in these patients. In addition, cancer, heart disease, and dementia were identified because of implications for end-of-life care, and atrial fibrillation was identified because of its increased risk of large cardioembolic strokes.

Palliative Care

Palliative care was identified using the ICD-9-CM procedure, code V66.7 (palliative care encounter [PCE]), in the hospital discharge data. This code is added by billing staff when components of palliative care, such as comfort care, end-of-life care, and hospice care, are mentioned in the treatment record of the patient and is independent of whether or not a palliative care specialist was consulted or not.²¹ The PCE code is not used for pain and symptom management. This article uses the term PCE to indicate the presence of V66.7 code in the patient's medical record. Several scenarios about the use of the V66.7 code in end-of-life and hospice care admissions and its interpretation by multiple national databases, such as CMS and US News and World Report, have been described.²² Recently, this code was examined in patients with ICH using NIS data from the previous decade.²³

Death

We used the healthcare cost and utilization project database uniform discharge disposition to track death during hospitalization. We compared the timing of death in PCE versus non-PCE patients. We defined early death as death occurring with hospital LOS ≤2 days. We explored implications of early death for stroke mortality as a CMS measure of high-quality care in the setting of PCE. As the healthcare

cost and utilization project database format changed in 2012, this combined analysis was limited to 2010 and 2011 data.

Statistical Analysis

Pearson χ^2 test was used to compare proportions between categories of PCE versus no PCE. Logistic regression was used to evaluate independent associations with PCE use. Covariates for logistic regression included age, race, sex, hospital characteristics, all-patient refined diagnosis-related group severity, and year. Statistical significance was defined as a *P* value of <0.05. Statistical analysis was performed using STATA data analysis and statistical software.

The all-patient refined diagnosis-related group, which assesses risk of mortality using an algorithm developed by 3 mol/L health information systems, was used to determine disease severity and its correlation with PCE. All-patient refined diagnosis-related group is a proprietary 4-point ordinal scale (minor, moderate, major, and extreme risk of mortality) derived from age, primary and secondary diagnoses, and procedures.²⁴

Results

We identified 395 411 adult patients with stroke. The majority of patients had ischemic strokes (86%) followed by ICH (10%) and SAH (4%). The mean age was 70.1 years (SD, 16), 52% were women and 69% were white. Among all patients with stroke, 24 641 (6.2%) received PCE, and this proportion increased with each study year from 5.4% in 2010 to 6.9% in 2012 (Table I in the online-only Data Supplement).

Palliative Care and Patient Characteristics

Bivariate analysis of pertinent variables is presented in Table 1. Although specific stroke severity scales (National Institutes of Health Stroke Scale, ICH score, Hunt/Hess) were not available in this cohort, proxies of overall illness severity, including the all-patient refined diagnosis-related group severity subclass and codes for intubation and coma, were associated with an increased rate of PCE use, whereas PEG placement was less common among patients with PCE (Table 1).

Using multivariate analysis, we found a variety of patient characteristics that were independently associated with the use of PCE (Table 2), including older age and female sex. Compared with whites, the rate of PCE use was significantly lower in blacks (odds ratio [OR], 0.62), Hispanics (OR, 0.67), and Asians (OR, 0.73). ICH, while representing only 10% of overall strokes, was associated with a higher rate of PCE use than ischemic stroke (OR, 3.40).

The mean LOS for all patients with stroke receiving PCE was 6.8 days (95% confidence interval, 6.66–6.87), which was significantly longer than in patients who did not receive PCE (5.7 days; 95% confidence interval, 5.64–5.69). When looking at each stroke subtype separately, this association was evident for patients with ischemic stroke (7.4 versus 6.2 days). Conversely, PCE was associated with shorter LOS in patients with ICH (5.0 versus 8.3 days) or SAH (6 versus 12 days; Table 3).

Palliative Care and Hospital Characteristics

Hospitals with higher PCE use included large hospitals (OR, 1.24 compared with small hospitals), urban teaching (OR, 1.1 compared with rural), nonprofit hospitals (OR, 1.22 compared with government hospitals), and western states (OR, 1.5

Table 1. Patient and Hospital Characteristics in Relation to Palliative Care Encounter (Bivariate Analysis)

Patient Characteristics	No Palliative Care, n (%)	Palliative Care, n (%)	P Value
Total	370 753 (93.8)	24 641 (6.2)	
Age, y			
<60	100 895 (97.0)	3132 (3.0)	<0.001
60–69	78 336 (95.9)	3351 (4.1)	<0.001
70–79	86 963 (93.9)	5675 (6.1)	<0.001
80+	104 575 (89.3)	12 484 (10.7)	<0.001
Sex			
Men	177 306 (94.8)	9770 (5.2)	<0.001
Women	193 394 (92.9)	14 870 (7.1)	<0.001
Race			
White	231 534 (93.0)	17 482 (7.0)	<0.001
Black	60 197 (96.2)	2364 (3.8)	<0.001
Hispanic	25 857 (95.4)	1254 (4.6)	<0.001
Asian/PI	8970 (93.6)	612 (6.4)	<0.001
Native American	1849 (94.2)	114 (5.8)	<0.001
Unknown	9632 (94.2)	590 (5.8)	<0.001
Stroke type			
Ischemic	322 959 (94.9)	17 221 (5.1)	<0.001
ICH	33 363 (84.7)	6032 (15.3)	<0.001
SAH	14 447 (91.2)	1389 (8.8)	<0.001
Insurance status			
Medicare	235 874 (93.0)	17 736 (7.0)	<0.001
Medicaid	29 399 (96.1)	1202 (3.9)	<0.001
Private insurance	74 194 (95.4)	3583 (4.6)	<0.001
Self-pay	19 989 (96.2)	788 (3.8)	<0.001
No charge	1779 (95.7)	81 (4.3)	<0.001
Other	8599 (88.5)	1121 (11.5)	<0.001
Hospital size			
Small	42 190 (95.0)	2201 (5.0)	<0.001
Medium	83 861 (94.1)	5303 (5.9)	<0.001
Large	241 268 (93.5)	16 852 (6.5)	<0.001
Hospital region			
Northeast	67 023 (94.2)	4104 (5.8)	<0.001
Midwest	83 178 (93.5)	5758 (6.5)	<0.001
South	152 744 (94.3)	9324 (5.7)	<0.001
West	67 824 (92.6)	5456 (7.4)	<0.001
Hospital type			
Rural	41 446 (94.9)	2227 (5.1)	<0.001
Urban nonteaching	142 453 (94.3)	8637 (5.7)	<0.001
Urban teaching	183 420 (93.2)	183 420 (6.8)	<0.001

(Continued)

Table 1. Continued

Patient Characteristics	No Palliative Care, n (%)	Palliative Care, n (%)	P Value
Hospital ownership			
Government (public)	44 922 (94.6)	2585 (5.4)	<0.001
Private (not for profit)	274 864 (93.3)	19 885 (6.7)	<0.001
Private (for profit)	47 533 (96.2)	1886 (3.8)	<0.001
Loss of function subclass*			
Minor	31 379 (98.4)	518 (1.6)	<0.001
Moderate	142 470 (97.7)	3369 (2.3)	<0.001
Major	129 756 (93.2)	9469 (6.8)	<0.001
Extreme	67 164 (85.6)	11 286 (14.4)	<0.001
Clinical characteristics			
Atrial fibrillation	81 191 (90.5)	8509 (9.5)	<0.001
Coronary artery disease	105 661 (93.2)	7760 (6.8)	<0.001
CHF	55 045 (91.2)	5291 (8.8)	<0.001
Dementia	5174 (93.1)	384 (6.9)	<0.001
Cancer	18 232 (92.5)	1468 (7.5)	<0.001
Comatose	4578 (62.1)	2799 (37.9)	<0.001
Inpatient procedures			
Mechanical ventilation	26 086 (82.2)	2799 (17.8)	<0.001
PEG placement	20 499 (94.9)	1113 (5.1)	<0.001
Tracheostomy	7045 (93.1)	520 (6.9)	0.020

APR-DRG indicates all-patient refined diagnosis-related group; CHF, congestive heart failure; ICH, intracerebral hemorrhage; PEG, percutaneous endoscopic gastrostomy; PI, Pacific Islander; and SAH, subarachnoid hemorrhage.

*Based on APR-DRG severity score.

compared with northeast hospitals). In general, hospitals with higher mortality also had higher use of PCE. However, this trend comes with a wide variability showing some hospitals with low PCE use and high mortality, as well hospitals with low mortality and high PCE use (Figure 1).

Palliative Care and Inpatient Mortality

Among all patients with stroke, 36 397 (9.2%) died in hospital, and the rate of death declined from 2010 (10.9%) to 2012 (9.8%; $P<0.001$; Table I in the online-only Data Supplement). Patient characteristics that were independently associated with higher mortality after stroke included older age (≥ 80 ; OR, 2.64 compared with <60), female sex (OR, 1.04), white race (OR, 0.77 for black versus white), ICH (OR, 4.76 compared with ischemic stroke), and non-Medicare insurance (OR self-pay 1.79 and private insurance 1.27 compared with Medicare). Hospital characteristics independently associated with higher mortality after stroke included small hospitals, hospitals in the northeast region, hospitals in rural areas, and public hospitals.

Among the patients who died, more than one third (38%) had received PCE (Table II in the online-only Data Supplement). The proportion of PCE was highest among patients dying with ICH (42%), followed by ischemic stroke (36%) and SAH (33%). Nonwhite races were less likely to

Table 2. Logistic Regression: Predictors of Palliative Care Encounter

Variable	Adjusted OR (95% CI)*	P Value
Discharge year		
2010	1.0 (reference)	
2011	1.17 (1.13–1.23)	<0.001
2012	1.42 (1.37–1.47)	<0.001
Age, y		
<60	1.0 (reference)	
60–69	1.77 (1.67–1.87)	<0.001
70–79	3.15 (2.97–3.34)	<0.001
80+	6.13 (5.79–6.49)	<0.001
Sex		
Men	1.0 (reference)	
Women	1.26 (1.22–1.30)	<0.001
Race		
White	1.0 (reference)	
Black	0.62 (0.59–0.65)	<0.001
Hispanic	0.67 (0.63–0.71)	<0.001
Asian/PI	0.73 (0.66–0.80)	<0.001
Native American	0.99 (0.80–1.21)	0.901
Unknown	0.76 (0.7–0.84)	<0.001
Stroke type		
Ischemic	1.0 (reference)	
ICH	3.40 (3.28–3.52)	<0.001
SAH	1.62 (1.52–1.74)	<0.001
Insurance status		
Medicare	1.0 (reference)	
Medicaid	1.32 (1.22–1.42)	<0.001
Private insurance	1.46 (1.39–1.53)	<0.001
Self-pay	1.70 (1.56–1.85)	<0.001
No charge	2.31 (1.81–2.96)	<0.001
Other	4.28 (3.96–4.62)	<0.001
Loss of function subclass		
Minor	1.0 (reference)	
Moderate	1.26 (1.14–1.39)	<0.001
Major	3.47 (3.16–3.81)	<0.001
Extreme	9.37 (8.52–10.3)	<0.001
Hospital size		
Small	1.0 (reference)	
Medium	1.18 (1.11–1.25)	<0.001
Large	1.24 (1.18–1.31)	<0.001
Hospital region		
Northeast	1.0 (reference)	
Midwest	1.23 (1.17–1.29)	<0.001

(Continued)

Table 2. Continued

Variable	Adjusted OR (95% CI)*	P Value
South	1.18 (1.14–1.23)	<0.001
West	1.50 (1.43–1.57)	<0.001
Hospital type		
Rural	1.0 (reference)	
Urban nonteaching	1.01 (0.96–1.06)	0.752
Urban teaching	1.10 (1.05–1.16)	<0.001
Hospital ownership		
Government (public)	1.0 (reference)	
Private (not for profit)	1.22 (1.17–1.28)	<0.001
Private (for profit)	0.65 (0.62–0.70)	<0.001

CI indicates confidence interval; ICH indicates intracerebral hemorrhage; OR, odds ratio; PI, Pacific Islander; and SAH, subarachnoid hemorrhage.

*OR calculated using multivariate logistic regression. All factors presented were included in the regression model except discharge status.

die in hospital, and among those who died, nonwhites were significantly less likely to receive PCE. The rates of death and PCE both increased with age, whereas the difference between the 2 decreased: among young patients (<60 years), more than twice as many patients died than received PCE. In the oldest patients (>90), more patients received PCE than died (Figure 2).

In the group of patients who died in hospital, the mean LOS was 7.0 days (95% confidence interval, 6.9–7.1). PCE use was associated with longer LOS in patients with ischemic stroke and with shorter LOS in patients with ICH and SAH. In decedents, PCE use was associated with a shorter LOS overall (6.2 versus 7.5 days) but with a longer LOS in patients with SAH (Table 3). In other words, PCE was associated with early death. The percentage of all PCE-related deaths was the highest in the earliest days of hospitalization both overall and for each stroke type (Figures 1 and 3).

Discussion

Using a well-established database of inpatient admissions in the United States, we found an overall rate of coding of PCEs among patients with stroke of 6.2% and an inpatient mortality rate of 9.2%. We observed substantial variation across patient and hospital characteristics and a strong correlation between palliative care use and death. Our findings have important implications for the use of hospital mortality rates as a CMS quality measure.

Palliative Care, Mortality, and Quality of Care Across Age, Race, and Sex

Consistent with the National Vital Statistics Report,¹ we observed that stroke admissions and in-hospital mortality increased with age. We found that the rate of PCE increased with age, and this difference was especially pronounced among decedents, where less than half of those younger than 60 years of age received PCE but all of those older than 90 years (Figure 2). In other words, older patients who die do so in the setting of PCE, whereas younger patients are more likely

Table 3. LOS With and Without PCE

Patient Characteristics	LOS Without PCE, d	LOS With PCE, d	Mean LOS, d
All patients	6.66	6.76	6.67
Ischemic	6.24*	7.40*	6.30*
ICH	8.33*	4.96*	7.81*
SAH	12.08*	6.46*	11.58*
In decedents	7.47*	6.16*	6.98*
Ischemic	9.16*	7.33*	8.50*
ICH	4.12	3.91	4.03
SAH	4.65*	5.57*	4.95*

ICH indicates intracerebral hemorrhage; LOS, length of stay; PCE, palliative care encounter; and SAH, subarachnoid hemorrhage.

*Values statistically significant with $P < 0.001$.

to die without PCE. This finding suggests a lower emphasis on life-prolonging care among older patients with stroke but may not indicate poor quality care. On the contrary, a recent study showed that the presence of DNR orders in patients with acute ischemic stroke, highly associated with older age and mortality, did not predict a lower incidence of stroke quality of care indicators.²⁵

Use of PCE and hospital mortality in the setting of PCE were also all more common in white people compared with blacks and Hispanics. This finding is consistent with a well-described racial variation in end-of-life care showing consistently lower rates of advance care planning,²⁶ DNR orders,²⁷ palliative care use²⁸ and end-of-life discussions,²⁹ and a higher rate of life-prolonging treatment, including PEG tube placement³⁰ among black patients with serious illness. Quality care indicators, however, are observed less frequently in black patients with stroke compared with white patients,³¹ and hospital deaths occur alongside high adherence to high-quality, evidence-based stroke care.¹⁶

Although not as clear, evidence for the association of female sex with palliative care use, as suggested in our study, has some support in the literature. Previous studies in patients with stroke have indicated higher rates of DNR orders and withdrawal of life-sustaining treatments in women and whites^{32–35} but slightly lower quality of care.^{36,37}




Figure 1. Rates of death and palliative care encounter (PCE) among patients with stroke by age.

Finally, our results suggest substantial practice variation of PCE use, consistent with the variation previously shown in end-of-life care after stroke, in particular in regards to the use of DNR orders,^{16,38} and prognostication.³⁹ The variation seen in our study may affect mortality in this group of patients, casting a shadow over the meaning and validity of mortality-based hospital comparisons that fail to account for PCE.

PCE (V66.7) Versus Palliative Care Services

The report card published by the center to advance palliative care showed that access to palliative care specialist services in US hospitals has increased in the past decade, but that a variability persists in regards to hospital size, location, and tax status.¹² In 2015, one third of US hospitals with 50 or more beds reported no palliative care services.¹² The reports' findings of a reduced rate of palliative care specialist services in smaller and nonacademic centers parallels the lower PCE rates in smaller, for-profit hospitals seen in our study, suggesting similar practice variations for specialist palliative care and palliative end-of-life care. Similarly, the geographic variation in palliative care specialist availability¹² corresponds with our finding of a higher PCE rate in the western states of the United States without substantial variation in mortality rates.

PCE and LOS

Among decedents, PCE was associated with a shorter LOS suggesting an earlier death through PCE and less days of aggressive life-sustaining treatment. LOS with PCE was longer in patients with ischemic stroke and shorter for ICH, which may be explained by a larger proportion of less-severe strokes on the one hand and later palliative care engagement for severe but nondeadly ischemic strokes on the other. This hypothesis was supported when we restricted the analysis to patients with ischemic stroke who died in the hospital: the trend reversed to shorter LOS with PCE. When looking only at patients with ICH, the association of PCE with shorter LOS was both seen in all patients with ICH and in decedents, possibly because of the high mortality and prognostic pessimism⁴⁰ in this stroke type. For the small group of patients with SAH, PCE was associated with shorter LOS, but here, the trend reversed when we looked only at patients who died. One possible explanation may be a difference in the culture of the medical services, given that SAH is typically managed by different medical teams than ischemic stroke and ICH.

Limitations

This study has several limitations, including those related to the retrospective analysis of the NIS database and the nature of an analysis based on ICD-9 coding. Large numbers in this data set lead to statistical significance even with small clinical changes. Owing to the nature of a preexisting database, important patient characteristics, such as stroke severity scales, are unavailable (eg, National Institutes of Health Stroke Scale). Second, ICD-9 coding is typically performed by the billing departments of hospitals based on language used by providers in their documentation. Provider documentation and billing guidelines may vary across individual departments, hospital types, geographic regions, and by individual administrative personnel. In addition,




Figure 2. Proportion of deaths per hospital length of stay among patients who received palliative care encounter (A). All patients (B through D) by stroke type. ICH indicates intracerebral hemorrhage; and SAH, subarachnoid hemorrhage.

because the V66.7 code is not linked to reimbursement, the documentation may be less reliable. It is possible that our observations indicate an increase in the coding of palliative care rather than an increase in the actual use of palliative care over time.




Figure 3. Hospital mortality and hospital palliative care encounter (PCE) use. Each dot represents one hospital. Hospitals with <10 stroke admissions or <10 palliative care encounters per year were excluded.

However, the patterns observed in this study correlate with other studies, suggesting a proportionate use of the PCE code. For example, the variability of PCEs across sampling year, age, region and hospital size, and ownership correlate with the availability of palliative care specialist services shown in the center to advance palliative care report card.¹³ Finally, documentation of PCE does not reflect the entirety of palliative care that a patient receives through primary or specialty palliative care. It also does not act as a surrogate for the degree to which goals of care, early comfort care measures, or surrogate decision making were addressed. Such services may be provided by the primary treating team without specific coding. More research is needed to build palliative and patient-centered care as a measurable healthcare quality metric.

Conclusions

Palliative care is increasing among patients with stroke, especially in larger hospitals. Disparities and variability in PCE and mortality across age, sex, race, region, and hospital characteristics are apparent. When evaluating 30-day mortality as a marker of quality of care, the presence or absence of PCE needs to be taken into account.

Disclosures

Dr Creutzfeldt receives support from the Cambia Health Foundation. The other authors report no conflicts.

References

- Melonie Heron PD. Deaths: leading causes for 2013. *NVSR*. 2016;65:1–95.
- Feigin VL, Lawes CM, Bennett DA, Barker-Collo SL, Parag V. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *Lancet Neurol*. 2009;8:355–369. doi: 10.1016/S1474-4422(09)70025-0.
- Creutzfeldt CJ, Longstreth WT, Holloway RG. Predicting decline and survival in severe acute brain injury: the fourth trajectory. *BMJ*. 2015;351:h3904. doi: 10.1136/bmj.h3904.
- van der Worp HB, Greving JP. Surviving space-occupying cerebral infarction: a fate worse than death? *Neurology*. 2010;75:676–677. doi: 10.1212/WNL.0b013e3181ee521.
- Creutzfeldt CJ, Holloway RG, Curtis JR. Palliative care: a core competency for stroke neurologists. *Stroke*. 2015;46:2714–2719. doi: 10.1161/STROKEAHA.115.008224.

6. Braun LT, Grady KL, Kutner JS, Adler E, Berlinger N, Boss R, et al; American Heart Association Advocacy Coordinating Committee. Palliative care and cardiovascular disease and stroke: a policy statement from the American Heart Association/American Stroke Association. *Circulation*. 2016;134:e198–e225. doi: 10.1161/CIR.0000000000000438.
7. Holloway RG, Arnold RM, Creutzfeldt CJ, Lewis EF, Lutz BJ, McCann RM, et al; American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, and Council on Clinical Cardiology. Palliative and end-of-life care in stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:1887–1916. doi: 10.1161/STR.0000000000000015.
8. Souter MJ, Blissitt PA, Blosser S, Bonomo J, Greer D, Jichici D, et al. Recommendations for the critical care management of devastating brain injury: prognostication, psychosocial, and ethical management: a position statement for healthcare professionals from the neurocritical care society. *Neurocrit Care*. 2015;23:4–13. doi: 10.1007/s12028-015-0137-6.
9. Lutz BJ, Green T. Nursing's role in addressing palliative care needs of stroke patients. *Stroke*. 2016;47:e263–e265. doi: 10.1161/STROKEAHA.116.013282.
10. Schutz RE, Coats HL, Engelberg RA, Curtis JR, Creutzfeldt CJ. Is there hope? Is she there? How families and clinicians experience severe acute brain injury. *J Palliat Med*. 2017;20:170–176. doi: 10.1089/jpm.2016.0286.
11. Liu Y, Kline D, Aerts S, Youngwerth JM, Kutner JS, Sillau S, et al. Inpatient palliative care for neurological disorders: lessons from a large retrospective series. *J Palliat Med*. 2017;20:59–64. doi: 10.1089/jpm.2016.0240.
12. America's Care of Serious Illness: 2015 State-by-state report card on access to palliative care in our nation's hospitals. Center to Advance Palliative Care Web site. <https://reportcard.capc.org>. Accessed July 1, 2015.
13. Dumanovsky T, Augustin R, Rogers M, Lettang K, Meier DE, Morrison RS. The growth of palliative care in U.S. hospitals: a status report. *J Palliat Med*. 2016;19:8–15. doi: 10.1089/jpm.2015.0351.
14. Measure Summary: Stroke: hospital 30-day, all-cause, risk-standardized mortality rate following acute ischemic stroke hospitalization. NQMC: 010156. Oct 2015. National Quality Measures Clearinghouse (NQMC). Agency for Healthcare Research and Quality Web site. <https://www.qualitymeasures.ahrq.gov>. Accessed July 1, 2016.
15. Hemphill JC III, Newman J, Zhao S, Johnston SC. Hospital usage of early do-not-resuscitate orders and outcome after intracerebral hemorrhage. *Stroke*. 2004;35:1130–1134. doi: 10.1161/01.STR.0000125858.71051.ca.
16. Kelly AG, Hoskins KD, Holloway RG. Early stroke mortality, patient preferences, and the withdrawal of care bias. *Neurology*. 2012;79:941–944. doi: 10.1212/WNL.0b013e3182666c40.
17. Introduction to the HCUP Nationwide Inpatient Sample (NIS), 2011. Rockville, MD: Agency for Healthcare Research and Quality Web site. Issued June 2013. Updated November 2015. http://www.hcup-us.ahrq.gov/db/nation/nis/NIS_Introduction_2011.jsp. Accessed July 1, 2015.
18. HCUP Nationwide Inpatient Sample (NIS) Overview. Healthcare Cost and Utilization Project (HCUP), 2010–2011. Agency for Healthcare Research and Quality Web site. <https://www.hcup-us.ahrq.gov/nisoverview.jsp>. Accessed July 1, 2015.
19. HCUP Nationwide Inpatient Sample (NIS) Overview. Healthcare Cost and Utilization Project (HCUP), 2012. Agency for Healthcare Research and Quality Web site. <https://www.hcup-us.ahrq.gov/nisoverview.jsp>. Accessed July 1, 2015.
20. Tirschwell DL, Longstreth WT Jr. Validating administrative data in stroke research. *Stroke*. 2002;33:2465–2470.
21. Brown F. *ICD-9-CM Coding Handbook, With Answers*. Revised Edition. Chicago, IL: AHA Press; 2012.
22. Cassel JB, Jones AB, Meier DE, Smith TJ, Spragens LH, Weissman D. Hospital mortality rates: how is palliative care taken into account? *J Pain Symptom Manage*. 2010;40:914–925. doi: 10.1016/j.jpainsymman.2010.07.005.
23. Murthy SB, Moradiya Y, Hanley DF, Ziai WC. Palliative care utilization in nontraumatic intracerebral hemorrhage in the United States. *Crit Care Med*. 2016;44:575–582. doi: 10.1097/CCM.0000000000001391.
24. Baram D, Daroowalla F, Garcia R, Zhang G, Chen JJ, Healy E, et al. Use of the all patient refined-diagnosis related group (APR-DRG) risk of mortality score as a severity adjustor in the medical ICU. *Clin Med Circ Respir Pulm Med*. 2008;2:19–25.
25. Reeves MJ, Myers LJ, Williams LS, Phipps MS, Bravata DM. Do-not-resuscitate orders, quality of care, and outcomes in veterans with acute ischemic stroke. *Neurology*. 2012;79:1990–1996. doi: 10.1212/WNL.0b013e3182735ced.
26. Narang AK, Wright AA, Nicholas LH. Trends in advance care planning in patients with cancer: results from a national longitudinal survey. *JAMA Oncol*. 2015;1:601–608. doi: 10.1001/jamaoncol.2015.1976.
27. Richardson DK, Zive DM, Newgard CD. End-of-life decision-making for patients admitted through the emergency department: hospital variability, patient demographics, and changes over time. *Acad Emerg Med*. 2013;20:381–387. doi: 10.1111/acem.12112.
28. Rush B, Hertz P, Bond A, McDermid RC, Celi LA. Use of palliative care in patients with end-stage COPD and receiving home oxygen: national trends and barriers to care in the United States. *Chest*. 2017;151:41–46. doi: 10.1016/j.chest.2016.06.023.
29. Mack JW, Paulk ME, Viswanath K, Prigerson HG. Racial disparities in the outcomes of communication on medical care received near death. *Arch Intern Med*. 2010;170:1533–1540. doi: 10.1001/archinternmed.2010.322.
30. George BP, Kelly AG, Schneider EB, Holloway RG. Current practices in feeding tube placement for US acute ischemic stroke inpatients. *Neurology*. 2014;83:874–882. doi: 10.1212/WNL.0000000000000764.
31. Fonarow GC, Reeves MJ, Zhao X, Olson DM, Smith EE, Saver JL, et al; Get With The Guidelines-Stroke Steering Committee and Investigators. Age-related differences in characteristics, performance measures, treatment trends, and outcomes in patients with ischemic stroke. *Circulation*. 2010;121:879–891. doi: 10.1161/CIRCULATIONAHA.109.892497.
32. Hemphill JC III. Do-not-resuscitate orders, unintended consequences, and the ripple effect. *Crit Care*. 2007;11:121. doi: 10.1186/cc5687.
33. Newman AJ III, Kvale EA, Williams BR, Bailey FA. What about the trach? Tracheotomy removal as a palliative care maneuver. *Am J Hosp Palliat Care*. 2007;24:371–375. doi: 10.1177/1049909107300214.
34. Qureshi AI, Adil MM, Suri MF. Rate of utilization and determinants of withdrawal of care in acute ischemic stroke treated with thrombolytics in USA. *Med Care*. 2013;51:1094–1100. doi: 10.1097/MLR.0b013e3182a95db4.
35. Zahuranec DB, Brown DL, Lisabeth LD, Gonzales NR, Longwell PJ, Smith MA, et al. Ethnic differences in do-not-resuscitate orders after intracerebral hemorrhage. *Crit Care Med*. 2009;37:2807–2811. doi: 10.1097/CCM.0b013e3181a56755.
36. McDermott M, Lisabeth LD, Baek J, Adelman EE, Garcia NM, Case E, et al. Sex disparity in stroke quality of care in a community-based study. *J Stroke Cerebrovasc Dis*. 2017;26:1781–1786. doi: 10.1016/j.jstrokecerebrovasdis.2017.04.006.
37. Reeves MJ, Fonarow GC, Zhao X, Smith EE, Schwamm LH; Get With The Guidelines-Stroke Steering Committee & Investigators. Quality of care in women with ischemic stroke in the GWTG program. *Stroke*. 2009;40:1127–1133. doi: 10.1161/STROKEAHA.108.543157.
38. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics-2016 update: a report from the American Heart Association. *Circulation*. 2016;133:e38–e360. doi: 10.1161/CIR.0000000000000350.
39. Zahuranec DB, Fagerlin A, Sánchez BN, Roney ME, Thompson BB, Fuhrel-Forbis A, et al. Variability in physician prognosis and recommendations after intracerebral hemorrhage. *Neurology*. 2016;86:1864–1871. doi: 10.1212/WNL.0000000000002676.
40. Creutzfeldt CJ, Becker KJ, Weinstein JR, Khot SP, McPharlin TO, Ton TG, et al. Do-not-attempt-resuscitation orders and prognostic models for intraparenchymal hemorrhage. *Crit Care Med*. 2011;39:158–162. doi: 10.1097/CCM.0b013e3181fb7b49.

Palliative Care for Hospitalized Patients With Stroke: Results From the 2010 to 2012 National Inpatient Sample

Tarvinder Singh, Steven R. Peters, David L. Tirschwell and Claire J. Creutzfeldt

Stroke. 2017;48:2534-2540; originally published online August 17, 2017;
doi: 10.1161/STROKEAHA.117.016893

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2017 American Heart Association, Inc. All rights reserved.
Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/48/9/2534>

Free via Open Access

Data Supplement (unedited) at:

<http://stroke.ahajournals.org/content/suppl/2017/08/17/STROKEAHA.117.016893.DC1>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Stroke* is online at:
<http://stroke.ahajournals.org/subscriptions/>

ONLINE SUPPLEMENT

Palliative care for hospitalized stroke patients: results from the 2010-2012 National Inpatient Sample

Supplemental Tables

TABLE I: Trends in Palliative care utilization.

Year	2010	2011	2012
Number of hospitals	1001	990	3829
Total number of stroke admissions	130,418	137,846	127,230
In-hospital deaths (% of stroke admissions)	12,773 (9.8%)	12,314 (8.9%)	11,310 (8.9%)
Total number of admissions receiving PCE	7,061	8,810	8,773
In-hospital stroke deaths receiving PCE (%)	32.7%	37.9%	42.0%
PCE per stroke admission (%)	5.4%	6.3%	6.9%

Table II: Discharge characteristics. All p < 0.01.

Discharge characteristics	No Palliative Care	Palliative Care
	n (%)	n (%)
Discharge Status (all patients, 2010-2012)		
Alive at Discharge	347,782 (93.87)	10,933 (44.39)
Died in Hospital	22,700 (6.13)	13,697 (55.61)
Discharge Status (only patients with discharge status, 2010-2011)		
Alive at Discharge except hospice	199,955 (90.05)	1,932 (14.16)
Died in Hospital	14,436 (6.50)	7,799 (57.20)
Discharged to hospice	7,648 (3.45)	3,904 (28.64)
Discharge Disposition (2010-2011)		
Routine	72,254 (32.54)	246 (1.80)
Skilled Nursing Facility (SNF)	44,218 (19.91)	1,198 (8.79)
Home Health Care	25,283 (11.39)	113 (0.83)
Rehab facility	39,462 (17.77)	79 (0.58)
Short-term hospital	6,631 (2.99)	69 (0.51)
Expired in hospital	14,376 (6.47)	7,666 (56.22)
Died in a medical facility	90 (0.04)	133 (0.09)
Hospice--Home	2,287 (1.03)	1,132 (8.30)
Hospice--Medical	5,361 (2.41)	2,772 (20.33)
Others	12,077 (5.45)	348 (2.55)