

## Implications of New High Blood Pressure Guidelines for Stroke Primary and Secondary Prevention

Daniel T. Lackland, DrPH

Since the report of the Society of Actuaries in the 1920s,<sup>1</sup> the excess disease risks of high blood pressure has been extensively studied, recognized, and documented. The linear relationship of blood pressure level and risk of stroke is evident among all adults regardless of age, sex, and race, leading to hypertension as the major global risk factor targeted for intervention.<sup>2-4</sup> The early clinical trials in the 1970s provided evidence that hypertension treatment and blood pressure reduction were associated with reduced stroke risks.<sup>5</sup> Epidemiology studies produced stroke risk estimates that were used in models to predict the impact of blood pressure reduction in the populations.<sup>6,7</sup> These estimates projected that specific reduction in blood pressures would correspond to reductions in population stroke rates (Table).

These risk estimates were shown to be valid as reductions in the distributions of population systolic blood pressures were detected from 1960 to 2008.<sup>8</sup> The greatest changes were detected for 60 to 74 year olds with median reductions from 148 to 129 mmHg and 90th percentile reductions from 188 to 156 mmHg. These lower blood pressure levels were most likely affected by hypertension treatment and management.<sup>9,10</sup> However, lower blood pressure levels were also identified in 20 to 29 year olds, suggesting that prevention and nonpharmacological strategies were effective as well.

Beginning in 1997 and continuing through 2017, the evidence from clinical trials and epidemiology studies was incorporated in a series of guidelines for detection, treatment, control, and prevention of high blood pressure.<sup>11-19</sup> With each rendition of the guidelines, refined strategies for hypertension intervention were developed by expert panels. In addition to blood pressure regimens and treatment protocols, hypertension

diagnostic procedures and algorithms are included in the recommendations. Although the reductions in stroke risks and population blood pressure distributions cannot be attributed to the recommendations from Guidelines, the implementation into practice has shown effects and impact.<sup>20,21</sup> These trends in population hypertension control and reduced stroke risks, along with the successful implementation of protocols such as demonstrated by Kaiser Permanente Northern California, would indicate the importance of maintaining intense recommendations in the guidelines. In contrast, an easing of the intensity of hypertension treatment and control may reverse the blood pressure-lowering and stroke risk reduction trends.

The reduced population stroke risk with reductions in blood pressure distributions has been proposed as the major public health achievements of the past 50 years.<sup>8</sup> As indicated, these trends are the result of the implementation of the evidence-based recommendations by healthcare providers and the adherence to the therapies by the population. It is expected that the intense recommendations of the 2017 Guidelines that builds on the evidence with new trial and study results will continue trends in the population cerebrovascular health.<sup>19,22</sup> Although these rates of stroke risk reduction and high blood pressure control are compelling, gaps in hypertension treatment and prevention remain, and it is critical to

- continue research to address the refinement of treatment protocols and the implementation;
- continue to critique recommendations and identify gaps in knowledge, behavior, and outcomes; and
- continue to form and maintain guideline panels charged with developing and enhancing evidence-based recommendations and protocols with the due diligence of the previous committees.

Still, it is important to recognize the remarkable trends in population blood pressure lowering and stroke risks reduction with the potential high impact from recommendations and the expectations with the new 2017 Guidelines.

### Disclosures

None.

### References

1. Society of Actuaries. *Build and Blood Pressure Study. Vol 1*. Chicago, IL: Society of Actuaries; 1959.
2. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R; Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903-1913. doi: 10.1016/S0140-6736(02)11911-8.
3. Lackland DT, Weber MA. Global burden of cardiovascular disease and stroke: hypertension at the core. *Can J Cardiol*. 2015;31:569-571. doi: 10.1016/j.cjca.2015.01.009.
4. Freis ED. The role of hypertension. *Am J Public Health*. 1960;50:11-13.
5. Freis ED. The Veterans Administration Cooperative Study on antihypertensive agents: implications for stroke prevention. *Stroke*. 1974;5:76-77.

**Table. Reduction in Population Systolic Blood Pressure Distributions With Projected Stroke Mortality Reduction**

Reduction in Systolic Blood Pressure, mmHg	Stroke Mortality Reduction, %
2	-6
3	-8
5	-14

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From the Department of Neurology, Medical University of South Carolina, Charleston.

Correspondence to Daniel T. Lackland, DrPH, Department of Neurology, Medical University of South Carolina, Harborview Office Tower, Suite 501, Charleston, SC 29425. E-mail lackland@musc.edu (*Stroke*. 2018;49:511-512).

DOI: 10.1161/STROKEAHA.118.020136.)

© 2018 American Heart Association, Inc.

*Stroke* is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.118.020136

6. Stamler R. Implications of the INTERSALT study. *Hypertension*. 1991;17(suppl 1):116–120.
7. Stamler J, Stamler R, Neaton JD, Wentworth D, Daviglius ML, Garside D, et al. Low risk-factor profile and long-term cardiovascular and noncardiovascular mortality and life expectancy: findings for 5 large cohorts of young adult and middle-aged men and women. *JAMA*. 1999;282:2012–2018.
8. Lackland DT, Roccella EJ, Deutsch AF, Fornage M, George MG, Howard G, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Quality of Care and Outcomes Research; Council on Functional Genomics and Translational Biology. Factors influencing the decline in stroke mortality: a statement from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:315–353. doi: 10.1161/01.str.0000437068.30550.cf.
9. Weber MA, Lackland DT. Hypertension: cardiovascular benefits of lowering blood pressure. *Nat Rev Nephrol*. 2016;12:202–204. doi: 10.1038/nrneph.2016.27.
10. Weber MA, Lackland DT. Contributions to hypertension public policy and clinical practice: a review of recent reports. *J Clin Hypertens (Greenwich)*. 2016;18:1063–1070. doi: 10.1111/jch.12902.
11. Report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure: a cooperative study. *JAMA*. 1977;237:255–261.
12. The 1980 report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure. *Arch Intern Med*. 1980;140:1280–1285.
13. The 1984 Report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure. *Arch Intern Med*. 1984;144:1045–1057.
14. The 1988 report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure. *Arch Intern Med*. 1988;148:1023–1038.
15. The fifth report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure (JNC V). *Arch Intern Med*. 1993;153:154–183.
16. The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med*. 1997;157:2413–2446.
17. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al; National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA*. 2003;289:2560–2572. doi: 10.1001/jama.289.19.2560.
18. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014;311:507–520. doi: 10.1001/jama.2013.284427.
19. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines [published online ahead of print November 13, 2017]. *Hypertension*. doi: 10.1161/HYP.0000000000000066. <http://hyper.ahajournals.org/content/hypertensionaha/early/2017/11/10/HYP.0000000000000065.full.pdf>. Accessed November 13, 2017.
20. Handler J, Lackland DT. Translation of hypertension treatment guidelines into practice: a review of implementation. *J Am Soc Hypertens*. 2011;5:197–207. doi: 10.1016/j.jash.2011.03.002.
21. Jaffe MG, Young JD. The Kaiser Permanente Northern California story: improving hypertension control from 44% to 90% in 13 years (2000–2013). *J Clin Hypertens (Greenwich)*. 2016;18:260–261.
22. Reboussin DM, Allen NB, Griswold ME, Guallar E, Hong Y, Lackland DT, et al. 2017 High Blood Pressure Systematic Review for the 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines [published online ahead of print November 13, 2017]. *Hypertension*. doi: 10.1161/HYP.0000000000000067. <http://hyper.ahajournals.org/content/hypertensionaha/early/2017/11/10/HYP.0000000000000067.full.pdf>. Accessed November 13, 2017.

KEY WORDS: Editorials ■ blood pressure ■ guideline ■ hypertension ■ risk ■ stroke

## Implications of New High Blood Pressure Guidelines for Stroke Primary and Secondary Prevention

Daniel T. Lackland

*Stroke*. 2018;49:511-512; originally published online February 12, 2018;  
doi: 10.1161/STROKEAHA.118.020136

*Stroke* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231  
Copyright © 2018 American Heart Association, Inc. All rights reserved.  
Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/49/3/511>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Stroke* is online at:  
<http://stroke.ahajournals.org/subscriptions/>