

## Rates of Spinal Cord Infarction After Repair of Aortic Aneurysm or Dissection

Gino Gialdini, MD; Neal S. Parikh, MD; Abhinaba Chatterjee, BS; Michael P. Lerario, MD; Hooman Kamel, MD; Darren B. Schneider, MD; Babak B. Navi, MD, MS; Santosh B. Murthy, MD, MPH; Costantino Iadecola, MD; Alexander E. Merkler, MD

**Background and Purpose**—The rate of spinal cord infarction (SCI) after surgical or endovascular repair of an aortic aneurysm or dissection is unclear.

**Methods**—Using administrative claims data, we identified adult patients discharged from nonfederal acute care hospitals in California, New York, and Florida who underwent surgical or endovascular repair of an aortic aneurysm or dissection between 2005 and 2013. Patients with SCI diagnosed before the aortic repair were excluded. Our primary outcome was an SCI during the index hospitalization for aortic repair. Descriptive statistics were used to estimate crude rates of SCI. Analyses were stratified by whether the aneurysm or dissection had ruptured and by type of repair (surgical versus endovascular).

**Results**—We identified 91 212 patients who had repair of an aortic aneurysm or dissection. SCI occurred in 235 cases (0.26%; 95% confidence interval [CI], 0.22%–0.29%). In patients with ruptured aneurysm or dissection, the rate of SCI was 0.74% (95% CI, 0.60%–0.88%) compared with 0.16% (95% CI, 0.13%–0.19%) with unruptured aneurysm. In secondary analyses, rates of SCI were similar after endovascular repair (0.91%; 95% CI, 0.62%–1.19%) compared with surgical repair (0.68%; 95% CI, 0.53%–0.83%;  $P=0.147$ ) of ruptured aortic aneurysm or dissection; however, rates of SCI were higher after surgical repair (0.20%; 95% CI, 0.15%–0.25%) versus endovascular repair (0.11%; 95% CI, 0.08%–0.14%;  $P<0.001$ ) of unruptured aneurysms.

**Conclusions**—SCI occurs in  $\approx 1$  in 130 patients undergoing aortic dissection or ruptured aortic aneurysm repair and in 1 in 600 patients undergoing unruptured aortic aneurysm repair. (*Stroke*. 2017;48:00–00. DOI: 10.1161/STROKEAHA.117.017071.)

**Key Words:** aortic aneurysm ■ endovascular procedures ■ hospitalization ■ infarction ■ spinal cord

Spinal cord infarction (SCI) accounts for up to 1% of all strokes<sup>1</sup> and can lead to severe disability, including paraplegia and tetraplegia. Both surgical and endovascular repair of an aortic aneurysm or dissection can lead to infarction of the spinal cord because the vascular supply of the spinal cord largely originates directly from the aorta.<sup>2</sup> The rate of SCI in patients undergoing aortic repair seems to vary depending on the presence and location of dissection,<sup>3,4</sup> aneurysm rupture,<sup>5</sup> and type of repair (surgical or endovascular).<sup>6,7</sup> Reported rates of SCI in these patients vary considerably, from 0.2% to 16%, and are largely based on single-center studies.<sup>3,8–12</sup> Furthermore, because new intraprocedural strategies to prevent SCI have been developed during the past several years, including cerebrospinal fluid drainage,<sup>13</sup> a better understanding of the risk of SCI after aortic repair is required. This is

particularly true because the number of patients who undergo repair of unruptured aneurysms in the United States is increasing.<sup>14</sup> We, therefore, sought to evaluate and compare the incidence of SCI after surgical and endovascular repair of aortic aneurysm and dissection in a large, heterogeneous cohort of patients from across the United States.

### Methods

#### Design

We performed a retrospective cohort study using administrative claims data from all inpatient discharges in California (2005–2011), Florida (2005–2013), and New York (2006–2013). Data from all hospitalizations at nonfederal hospitals were collected by the California Office of Statewide Health Planning and Development, the New York Statewide Planning and Research Cooperative System, and

Received February 14, 2017; final revision received May 3, 2017; accepted June 5, 2017.

From the Clinical and Translational Neuroscience Unit, Feil Family Brain and Mind Research Institute (G.G., N.S.P., A.C., H.K., B.B.N., S.B.M., C.I., A.E.M.), Department of Neurology (N.S.P., M.P.L., H.K., B.B.N., S.B.M., C.I., A.E.M.), and Department of Vascular and Endovascular Surgery (D.B.S.), Weill Cornell Medicine, New York, NY; and Department of Neurology, New York-Presbyterian Queens, Flushing (M.P.L.).

Presented in part at the International Stroke Conference, Houston, TX, February 23, 2017, and the American Academy of Neurology, Boston, MA, April 28, 2017.

Jeffrey Saver, MD, was the Consulting Editor for this paper.

Correspondence to Alexander E. Merkler, MD, Clinical and Translational Neuroscience Unit, Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, 525 E, 68th St, F610, New York, NY 10065. E-mail alm9097@med.cornell.edu

© 2017 American Heart Association, Inc.

*Stroke* is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.117.017071

the Florida Agency for Health Care Administration. These databases were deidentified by the Agency for Healthcare Research and Quality, which assigned patients a personal linkage number that allows them to be anonymously tracked across multiple hospitalizations. Up to 25 discharge diagnoses and 21 procedures were coded for each encounter using the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)* system with each diagnosis labeled as having been present before hospital admission or having newly developed during the hospitalization. Because of the deidentified nature of the data, the study was exempt from formal review by the institutional review board at Weill Cornell Medicine.

## Patients

We included all patients at the time of their first recorded hospitalization for a surgical or endovascular repair of a thoracic, thoracoabdominal, or abdominal aortic dissection or aneurysm. Patients undergoing repair for both ruptured and unruptured aneurysms were included. Cases were identified using *ICD-9-CM* procedure codes for endovascular (39.71 and 39.73) or surgical aortic repair (38.34, 38.44-45, 39.23-26, 39.57-58) in addition to diagnosis codes for aortic dissection or aneurysm (*ICD-9-CM* codes 441.0-9) in any discharge diagnosis position within the same encounter. This approach has been used previously.<sup>15,16</sup>

## Measurements

Our primary outcome was SCI defined by the *ICD-9-CM* code 336.1 in any discharge diagnosis position during the index hospitalization for repair of aortic aneurysm or dissection. We excluded cases with present-on-admission codes for SCI and any cases with a prior history of SCI before index hospitalization for aortic repair. In a secondary analysis, we expanded our definition of SCI to include any new-onset motor impairment coded as tetraplegia, paraplegia, and diplegia (*ICD-9-CM* codes 344.0-2).

Using *ICD-9-CM* codes, we identified the following vascular risk factors in our cohort: hypertension, diabetes mellitus, coronary artery disease, atrial fibrillation, peripheral vascular disease, congestive heart failure, chronic kidney disease, tobacco use, alcohol abuse, and chronic obstructive pulmonary disease. In addition, we included the Elixhauser comorbidity index, a comprehensive set of comorbidity measures commonly used for administrative data, to address disease severity.<sup>17</sup>

## Statistical Analysis

We compared patients' baseline characteristics using the  $\chi^2$  test for categorical variables and the Mann-Whitney *U* test for continuous variables because the data were not normally distributed. Exact confidence intervals (CIs) were used to report crude rates of SCI in patients who underwent a surgical or endovascular aortic repair. We evaluated the rates of SCI in patients who had an emergency surgery for either a ruptured aneurysm or aortic dissection and then separately in patients who had nonemergent surgery for an unruptured aneurysm. We further stratified our results for these 2 groups by treatment approach: surgical versus endovascular. Because patients who have a ruptured aortic aneurysm or dissection have high mortality,<sup>16</sup> and in particular, intraoperative mortality, we included only patients who survived to the end of the index hospitalization in a sensitivity analysis using the same statistical tests as described above. All statistical analyses were performed by G.G. and A.E.M. using Stata/MP (version 13, College Station, TX). The threshold of statistical significance was set at  $\alpha=0.05$ .

## Results

We identified 91 212 patients who underwent aortic aneurysm or dissection repair, 38 432 of whom received surgical intervention and 51 427 of whom received endovascular intervention. Of these, 1353 patients (1.48%) underwent both surgical and endovascular repair procedures within the same

encounter. A ruptured aortic aneurysm or dissection was diagnosed in 15 304 patients, whereas 74 555 patients presented with unruptured aneurysms. SCI was identified in 235 cases (0.26%; 95% CI, 0.22%–0.29%) and was more common among patients who were younger, women, non-white, and in patients with a higher burden of vascular risk factors (Table 1). SCI occurred more often in patients who underwent repair of a ruptured aortic aneurysm or aortic dissection (114 cases, 0.74%; 95% CI, 0.60%–0.88%) as compared with in patients who underwent repair of an unruptured aneurysm (106 cases, 0.16%; 95% CI, 0.13%–0.19%).

In patients with aortic dissection or ruptured aortic aneurysm, there was no difference in the rate of SCI after surgical repair (76 cases, 0.68%; 95% CI, 0.53%–0.83%) as compared

**Table 1. Characteristics of Patients, Stratified by Spinal Cord Infarction**

Characteristic*	SCI (n=235)	No SCI (n=90 977)	P Value
Age, mean (SD), y	69.1 (11.7)	71.1 (11.3)	0.008
Female	87 (37.0)	21 871 (24.0)	<0.001
Race†			
White	172 (74.8)	72 545 (81.0)	0.06
Black	21 (9.1)	4656 (5.2)	
Hispanic	21 (9.1)	6560 (7.3)	
Asian/other	16 (7.0)	5835 (6.5)	
Payment source‡			
Medicare	165 (69.8)	65 684 (72.2)	0.002
Medicaid	16 (7.2)	3482 (3.8)	
Private	41 (17.4)	18 901 (20.8)	
Self-pay/other	13 (5.6)	2906 (3.2)	
Hypertension	179 (76.2)	68 031 (74.8)	0.60
Diabetes mellitus	49 (20.9)	18 252 (20.1)	0.78
Coronary heart disease	80 (34.0)	40 478 (44.5)	0.001
Congestive heart failure	34 (14.5)	11 357 (12.5)	0.36
Peripheral vascular disease	94 (40.0)	22 058 (26.8)	<0.001
Chronic obstructive pulmonary disease	75 (31.9)	24 388 (26.8)	0.08
Chronic kidney disease	61 (26.0)	11 054 (12.2)	<0.001
Atrial fibrillation	65 (27.7)	18 920 (20.8)	0.01
Tobacco use	32 (13.6)	21 759 (23.9)	<0.001
Alcohol use	45 (19.2)	18 307 (20.1)	0.71
Elixhauser comorbidities§, mean (SD)	4.3 (1.9)	2.7 (1.6)	<0.001

SCI indicates spinal cord infarction.

\*Data are presented as number (%) unless otherwise specified.

†Self-reported by patients or their surrogates. Numbers do not sum to group totals because of missing race/ethnicity data in 2.2% of patients.

‡Numbers do not sum to group totals because of missing payment-source data in <0.01% of patients.

§Numbers represent the number of Elixhauser comorbid conditions, which comprise a comprehensive set of 28 comorbidity measures for use with large administrative datasets.

with endovascular repair (38 cases, 0.91%; 95% CI, 0.62%–1.19%;  $P=0.147$ ). However, in patients undergoing repair of an unruptured aneurysm, the rate of SCI appeared higher after surgical therapy (55 cases, 0.20%; 95% CI, 0.15%–0.25%) than after endovascular therapy (51 cases, 0.11%; 95% CI, 0.08%–0.14%;  $P<0.001$ ). Characteristics of patients undergoing surgical or endovascular repair are summarized in Tables 2 and 3. Finally, of the 1353 patients who had codes for both

surgical and endovascular aortic repair during the index hospitalization, 15 developed an SCI (1.11%; 95% CI, 0.55%–1.67%). In a sensitivity analysis after excluding patients who died during the hospitalization for aortic repair, the overall SCI rate was similar (0.21%; 95% CI, 0.18%–0.24%) as were the rates of SCI stratified by treatment approach. When we included new paraplegia, diplegia, and tetraplegia within the definition of SCI, the overall rate of SCI was 0.56% (95%

**Table 2. Characteristics of Patients With Ruptured Aortic Aneurysm or Dissection, Stratified by Type of Aortic Repair\***

Characteristic†	Endovascular (n=4179)	Surgery (n=11 125)	P Value
SCI	38 (0.9)	76 (0.7)	0.15
Age, mean (SD), y	70.7 (13.2)	65.8 (13.9)	0.02
Female	1220 (29.2)	3332 (29.9)	0.35
Race‡			
White	2883 (70.6)	7635 (70.6)	0.06
Black	480 (11.8)	1180 (10.9)	
Hispanic	330 (8.0)	1026 (9.5)	
Asian	194 (4.8)	477 (4.4)	
Other	194 (4.8)	492 (4.6)	
Payment source§			
Medicare	2822 (67.5)	6202 (55.7)	<0.001
Medicaid	286 (6.8)	1034 (9.3)	
Private	793 (19.0)	3000 (27.0)	
Self-pay	125 (3.0)	461 (4.2)	
Other	153 (3.7)	427 (3.8)	
Hypertension	3201 (76.6)	7949 (71.5)	<0.001
Diabetes mellitus	908 (21.7)	2135 (19.2)	<0.001
Coronary heart disease	1514 (36.2)	3581 (32.2)	<0.001
Congestive heart failure	611 (14.6)	2094 (18.8)	<0.001
Peripheral vascular disease	994 (23.8)	2163 (19.4)	<0.001
Chronic obstructive pulmonary disease	1008 (24.1)	2377 (21.4)	<0.001
Chronic kidney disease	731 (17.5)	1577 (14.2)	<0.001
Atrial fibrillation	862 (20.6)	3349 (30.1)	<0.001
Tobacco use	738 (17.7)	1329 (11.9)	<0.001
Alcohol use	856 (20.5)	2512 (22.6)	0.005
Elixhauser comorbidities,   mean (SD)	3.1 (1.7)	3.1 (1.8)	0.02

SCI indicates spinal cord infarction.

\*Data do not include cases who underwent combined surgical plus endovascular procedures.

†Data are presented as number (%) unless otherwise specified.

‡Self-reported by patients or their surrogates. Numbers do not sum to group totals because of missing race/ethnicity data in 2.8% of patients.

§Numbers do not sum to group totals because of missing payment-source data in <0.01% of patients.

||Numbers represent the number of Elixhauser comorbid conditions, which comprise a comprehensive set of 28 comorbidity measures for use with large administrative datasets.

**Table 3. Characteristics of Patients With Unruptured Aneurysm, Stratified by Type of Aortic Repair\***

Characteristic†	Endovascular (n=47 248)	Surgery (n=27 307)	P Value
SCI	51 (0.1)	55 (0.2)	<0.001
Age, mean (SD), y	74.4 (8.7)	67.5 (11.9)	<0.001
Female	8867 (18.8)	8041 (29.5)	<0.001
Race‡			
White	39 345 (84.5)	21 669 (80.8)	<0.001
Black	1786 (3.8)	1119 (4.2)	
Hispanic	2967 (6.4)	2140 (8.0)	
Asian	1171 (2.5)	817 (3.0)	
Other	1322 (2.8)	1088 (4.0)	
Payment source§			
Medicare	38 719 (82.0)	17 056 (62.5)	<0.001
Medicaid	923 (2.0)	1197 (4.4)	
Private	6802 (14.3)	8134 (29.8)	
Self-pay	229 (0.5)	294 (1.1)	
Other	574 (1.2)	624 (2.2)	
Hypertension	36 014 (76.2)	20 028 (73.3)	<0.001
Diabetes mellitus	9441 (20.0)	5530 (20.3)	0.40
Coronary heart disease	22 849 (48.4)	12 013 (44.0)	<0.001
Congestive heart failure	4039 (8.6)	4477 (16.4)	<0.001
Peripheral vascular disease	11 427 (24.2)	6900 (25.3)	0.001
Chronic obstructive pulmonary disease	13 024 (27.6)	7577 (27.8)	0.60
Chronic kidney disease	5671 (12.0)	2939 (10.8)	<0.001
Atrial fibrillation	6850 (14.5)	7638 (28.0)	<0.001
Tobacco use	13 240 (28.0)	6164 (22.6)	<0.001
Alcohol use	9031 (19.1)	5652 (20.7)	<0.001
Elixhauser comorbidities,   mean (SD)	2.4 (1.5)	3.0 (1.7)	<0.001

SCI indicates spinal cord infarction.

\*Data do not include cases who underwent combined surgical plus endovascular procedures.

†Data are presented as number (%) unless otherwise specified.

‡Self-reported by patients or their surrogates. Numbers do not sum to group totals because of missing race/ethnicity data in 1.8% of patients.

§Numbers do not sum to group totals because of missing payment-source data in <0.01% of patients.

||Numbers represent the number of Elixhauser comorbid conditions, which comprise a comprehensive set of 28 comorbidity measures for use with large administrative datasets.

CI, 0.51%–0.61%), and again, the outcomes were unchanged stratified by treatment approach.

## Discussion

In a large, heterogeneous cohort of patients who underwent surgical or endovascular repair of an aortic dissection or aneurysm, we found that the overall rate of SCI was 0.26%. The rate of SCI was significantly higher (1 in 130 cases) in patients who were treated for a ruptured aortic aneurysm or dissection as compared with an unruptured aneurysm (1 in 600). For patients undergoing treatment of ruptured aortic aneurysm or dissection, the rate of SCI was similar after endovascular compared with surgical repair. However, for patients with unruptured aneurysm, surgical therapy was associated with a higher risk of SCI compared with endovascular repair.

Our findings are consistent with and build on prior analyses. The overall rate of SCI in our data seems similar to what was reported by Qureshi et al<sup>18</sup> in a large, population-based study of administrative claims data, in which the overall rate of SCI after endovascular stent graft placement for aortic aneurysm or dissection repair was 0.2% in patients admitted between 2002 and 2011. However, these authors included only patients who underwent endovascular aortic repair and did not separately examine patients undergoing repair of ruptured versus unruptured aneurysms. In the only prospective trial evaluating complications after surgical or endovascular treatment of ruptured abdominal aneurysms, the rate of SCI was 0.9%<sup>19</sup>; however, the number of patients enrolled in the trial was low (116) and there was only 1 outcome. Furthermore, this trial was performed in academic hospitals in the Netherlands and therefore may not be generalizable to patients undergoing aortic repair in the United States.

In contrast to prior studies, we were able to compare rates of in-hospital SCI in patients undergoing repair of ruptured aortic aneurysms and dissection versus those undergoing repair of unruptured aneurysms. The relatively low rate of SCI seen after repair of unruptured aortic aneurysm is reassuring because these patients are presumably neurologically normal before surgery. In addition, our findings are consistent with prior reports of low rates of SCI after elective open thoracic aneurysm repair.<sup>20</sup> We did not find any differences according to treatment approach (surgery versus endovascular repair) in patients with ruptured aortic aneurysm or aortic dissection. However, we found a higher rate of SCI after surgical repair in patients with unruptured aneurysm or dissection. Our findings, if confirmed, may be relevant to treatment approach selection for elective repair. Irrespective of treatment approach, because the number of patients undergoing repair of unruptured aortic aneurysms is increasing in frequency in the United States,<sup>14</sup> further research to prevent ischemic complications after aortic aneurysm repair is warranted.<sup>21</sup>

The results of our study should be considered in the context of several limitations. First, We defined SCI through the use of *ICD-9-CM* code 336.1 in any diagnostic position. This diagnosis algorithm has been previously used in a study analyzing claims-based data,<sup>18</sup> but its sensitivity and specificity have not been conclusively established. However, *ICD-9-CM* hospital discharge codes have been shown to be generally reliable for major diseases.<sup>22</sup> To try to capture all cases of SCI, in a sensitivity analysis we also included patients with diagnoses

of tetraplegia, diplegia, or paraplegia. Second, because a limited number of SCI occurred in the non-white population, our results may not be generalizable to non-white patients undergoing aortic repair. Third, we lacked data on spinal cord imaging which are the most accurate, albeit limited test, to confirm SCI.<sup>23</sup> Fourth, because of the nature of the data, we could not differentiate whether SCI occurred as direct sequela of a ruptured aneurysm or aortic dissection or due to complications of vessel repair; however, we excluded cases where SCI was present on admission. Fifth, we may have missed cases of SCI in patients who died perioperatively and therefore were not diagnosed with SCI. However, our results were similar in a sensitivity analysis excluding patients who died during the index hospitalization for aortic repair. Sixth, because of the administrative nature of the data, we lacked important granular data that may have affected the risk of perioperative SCI, including aneurysm size at the time of repair, aneurysm growth over time, duration of procedure, and length of aorta replaced or repaired. Furthermore, we were unable to account for collateral blood supply to the spinal cord or use of cerebrospinal fluid drainage, which may affect the risk of SCI after aortic repair.<sup>13,24</sup>

## Conclusion

In a large, population-based cohort of patients, we found that SCI occurs in ≈1 in 130 patients who undergo a surgical or endovascular repair of an aortic dissection or ruptured aneurysm repair and 1 in 600 patients who undergo a surgical or endovascular repair of an unruptured aortic aneurysm. Given the relative frequency of this devastating complication of aortic repair, further study into the optimal method to prevent perioperative SCI is recommended.

## Acknowledgments

We are grateful to Monica Chen for her copyediting and clerical assistance.

## Sources of Funding

Dr Gialdini is supported by the Feil Family Foundation. Dr Murthy is supported by the American Brain Foundation and the American Academy of Neurology. Dr Kamel is supported by National Institutes of Health (NIH) grants K23NS082367 and R01NS097443, as well as the Michael Goldberg Research Fund. Dr Navi is supported by NIH grant K23NS091395 and the Florence Gould Endowment for Discovery in Stroke. Dr Iadecola is supported by NIH grants R37NS089323, R01NS034179, R01NS095441, and R01NS100447.

## Disclosures

None.

## References

- Romi F, Naess H. Spinal cord infarction in clinical neurology: a review of characteristics and long-term prognosis in comparison to cerebral infarction. *Eur Neurol*. 2016;76:95–98. doi: 10.1159/000446700.
- Colman MW, Hornicek FJ, Schwab JH. Spinal cord blood supply and its surgical implications. *J Am Acad Orthop Surg*. 2015;23:581–591. doi: 10.5435/JAAOS-D-14-00219.
- Gaul C, Dietrich W, Friedrich I, Sirch J, Erbguth FJ. Neurological symptoms in type A aortic dissections. *Stroke*. 2007;38:292–297. doi: 10.1161/01.STR.0000254594.33408.b1.
- Zhang L, Sun XG, Yu CT, Chang Q, Qian XY. Intercostal artery reconstruction: the simple and effective technique on spinal cord

- protection during thoracoabdominal aortic replacement. *Ann Vasc Surg.* 2016;34:62–67. doi: 10.1016/j.avsg.2015.12.030.
5. Coselli JS, LeMaire SA, Preventza O, de la Cruz KI, Cooley DA, Price MD, et al. Outcomes of 3309 thoracoabdominal aortic aneurysm repairs. *J Thorac Cardiovasc Surg.* 2016;151:1323–1337. doi: 10.1016/j.jtcvs.2015.12.050.
  6. Greenberg RK, Lu Q, Roselli EE, Svensson LG, Moon MC, Hernandez AV, et al. Contemporary analysis of descending thoracic and thoracoabdominal aneurysm repair: a comparison of endovascular and open techniques. *Circulation.* 2008;118:808–817. doi: 10.1161/CIRCULATIONAHA.108.769695.
  7. Badger S, Bedenis R, Blair PH, Ellis P, Kee F, Harkin DW. Endovascular treatment for ruptured abdominal aortic aneurysm. *Cochrane Database Syst Rev.* 2014;CD005261. doi: 10.1002/14651858.CD005261.pub3.
  8. Bisdas T, Panuccio G, Sugimoto M, Torsello G, Austermann M. Risk factors for spinal cord ischemia after endovascular repair of thoracoabdominal aortic aneurysms. *J Vasc Surg.* 2015;61:1408–1416. doi: 10.1016/j.jvs.2015.01.044.
  9. Ma WG, Zhang W, Wang LF, Zheng J, Ziganshin BA, Charilaou P, et al. Type A aortic dissection with arch entry tear: surgical experience in 104 patients over a 12-year period. *J Thorac Cardiovasc Surg.* 2016;151:1581–1592. doi: 10.1016/j.jtcvs.2015.11.056.
  10. Afifi RO, Sandhu HK, Leake SS, Boutros ML, Kumar V 3rd, Azizzadeh A, et al. Outcomes of patients with acute type B (DeBakey III) aortic dissection: a 13-year, single-center experience. *Circulation.* 2015;132:748–754. doi: 10.1161/CIRCULATIONAHA.115.015302.
  11. Lee SJ, Kim JH, Na CY, Oh SS, Kim YM, Lee CK, et al. Eleven years of experience with the neurologic complications in Korean patients with acute aortic dissection: a retrospective study. *BMC Neurol.* 2013;13:46. doi: 10.1186/1471-2377-13-46.
  12. Ullery BW, Cheung AT, Fairman RM, Jackson BM, Woo EY, Bavaria J, et al. Risk factors, outcomes, and clinical manifestations of spinal cord ischemia following thoracic endovascular aortic repair. *J Vasc Surg.* 2011;54:677–684. doi: 10.1016/j.jvs.2011.03.259.
  13. Uchida N. How to prevent spinal cord injury during endovascular repair of thoracic aortic disease. *Gen Thorac Cardiovasc Surg.* 2014;62:391–397. doi: 10.1007/s11748-014-0395-9.
  14. Karthikesalingam A, Vidal-Diez A, Holt PJ, Loftus IM, Schermerhorn ML, Soden PA, et al. Thresholds for abdominal aortic aneurysm repair in England and the United States. *N Engl J Med.* 2016;375:2051–2059. doi: 10.1056/NEJMoa1600931.
  15. Edwards ST, Schermerhorn ML, O'Malley AJ, Bensley RP, Hurks R, Cotterill P, et al. Comparative effectiveness of endovascular versus open repair of ruptured abdominal aortic aneurysm in the Medicare population. *J Vasc Surg.* 2014;59:575–582. doi: 10.1016/j.jvs.2013.08.093.
  16. Karthikesalingam A, Holt PJ, Vidal-Diez A, Ozdemir BA, Poloniecki JD, Hinchliffe RJ, et al. Mortality from ruptured abdominal aortic aneurysms: clinical lessons from a comparison of outcomes in England and the USA. *Lancet.* 2014;383:963–969. doi: 10.1016/S0140-6736(14)60109-4.
  17. Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Med Care.* 1998;36:8–27.
  18. Qureshi AI, Chughtai M, Malik AA. Analysis of spinal cord infarction associated with aortic stent graft placement using nationwide inpatient sample (2002–2011). *J Vasc Interv Neurol.* 2016;8:9–16.
  19. Reimerink JJ, Hoornweg LL, Vahl AC, Wisselink W, van den Broek TA, Legemate DA, et al; Amsterdam Acute Aneurysm Trial Collaborators. Endovascular repair versus open repair of ruptured abdominal aortic aneurysms: a multicenter randomized controlled trial. *Ann Surg.* 2013;258:248–256. doi: 10.1097/SLA.0b013e31828d4b76.
  20. Murphy EH, Stanley GA, Arko MZ, Davis CM 3rd, Modrall JG, Arko FR 3rd. Effect of ethnicity and insurance type on the outcome of open thoracic aortic aneurysm repair. *Ann Vasc Surg.* 2013;27:699–707. doi: 10.1016/j.avsg.2012.08.011.
  21. Salzler GG, Meltzer AJ, Mao J, Isaacs A, Connolly PH, Schneider DB, et al. Characterizing the evolution of perioperative outcomes and costs of endovascular abdominal aortic aneurysm repair. *J Vasc Surg.* 2015;62:1134–1139. doi: 10.1016/j.jvs.2015.06.138.
  22. Quan H, Parsons GA, Ghali WA. Assessing accuracy of diagnosis-type indicators for flagging complications in administrative data. *J Clin Epidemiol.* 2004;57:366–372. doi: 10.1016/j.jclinepi.2003.01.002.
  23. Novy J, Carruzzo A, Maeder P, Bogousslavsky J. Spinal cord ischemia: clinical and imaging patterns, pathogenesis, and outcomes in 27 patients. *Arch Neurol.* 2006;63:1113–1120. doi: 10.1001/archneur.63.8.1113.
  24. Banga PV, Oderich GS, Reis de Souza L, Hofer J, Cazares Gonzalez ML, Pulido JN, et al. Neuromonitoring, cerebrospinal fluid drainage, and selective use of iliofemoral conduits to minimize risk of spinal cord injury during complex endovascular aortic repair. *J Endovasc Ther.* 2016;23:139–149. doi: 10.1177/15266602815620898.

# Stroke



# Stroke

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## **Rates of Spinal Cord Infarction After Repair of Aortic Aneurysm or Dissection** Gino Gialdini, Neal S. Parikh, Abhinaba Chatterjee, Michael P. Lerario, Hooman Kamel, Darren B. Schneider, Babak B. Navi, Santosh B. Murthy, Costantino Iadecola and Alexander E. Merkler

*Stroke*. published online June 27, 2017;  
*Stroke* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231  
Copyright © 2017 American Heart Association, Inc. All rights reserved.  
Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/early/2017/06/27/STROKEAHA.117.017071>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Stroke* is online at:  
<http://stroke.ahajournals.org/subscriptions/>