

Stroke Risk Factors Unique to Women

Stacie L. Demel, DO, PhD; Steven Kittner, MD, MPH; Sylvia H. Ley, PhD, RD;
Mollie McDermott, MD, MS; Kathryn M. Rexrode, MD, MPH

Stroke is the third leading cause of death in women in the United States and is a leading cause of disability. Each year 55 000 more women than men have a stroke, a discrepancy largely driven by longer life expectancy in women (www.stroke.org). Although the majority of stroke incidence can be attributed to traditional vascular risk factors that occur in both men and women, including hypertension, hyperlipidemia, diabetes mellitus, smoking, and atrial fibrillation, there are several stroke risk factors that are specific to women. Specifically, differences in sex hormones, exogenous estrogens, and pregnancy exposures are factors exclusively experienced by women. In this review, we will summarize the current state of the literature with regards to women-specific factors, such as endogenous hormone levels, exogenous hormone therapy, pregnancy, parity, timing of age at menarche, and menopause in relation to stroke risk.

Methods

The following terms were searched with women and stroke in PubMed and Google Scholar, mainly for original articles and meta-analysis/systematic reviews: estrogens, estradiol, testosterone, DHEAS, menarche, menopause, oophorectomy, postmenopausal hormone therapy, oral contraception, transgender, transmen, transwomen, pregnancy, peripartum, postpartum, and parity. The following search was also performed: therapy OR treatment OR secondary prevention AND pregnancy AND stroke. The resultant literature was reviewed by the authors, and the data covering the topics outlined below were reviewed in this manuscript.

Endogenous Estrogen State

Endogenous Hormone Levels

Data on the relationship of endogenous sex hormones and risk of stroke in women are relatively limited. Estrogen levels fluctuate dramatically in women with the menstrual cycle and then drop dramatically in the menopausal transition and in postmenopause. In data from the Copenhagen City Study, neither high nor low estradiol levels were associated with increased risk of ischemic stroke.¹ In premenopausal women, those in the lowest 10th percentile of estradiol had a >2-fold increased risk of ischemic stroke, but this was based on small case

numbers. There was no relationship observed for postmenopausal estradiol levels and risk of ischemic stroke. Similarly, the authors of a study of French women over the age of 65 years found no association between estradiol levels and risk of ischemic stroke.² In the Study of Osteoporotic Fractures, women in the highest category of free estrogen index had a higher age-adjusted risk of ischemic stroke, but this was not independent of standard stroke risk factors, including hypertension, diabetes mellitus, and adiposity.³ Subsequently, a meta-analysis of the 3 available studies found no association, further supporting the lack of a relationship between estradiol levels and risk of ischemic stroke.¹

Testosterone levels are more stable than estrogen levels across the lifespan in women, with relatively constant levels from ages 30 to 70 years.¹ Although low testosterone levels have been associated with increased stroke risk in men, no clear relationship has been seen for testosterone levels and risk of stroke in women.¹ The investigators of the aforementioned study of French women over the age of 65 years found no association between high or low testosterone levels and risk of stroke.²

Dehydroepiandrosterone, an adrenal hormone which can also be used for the synthesis of estrogen and testosterone, has also been investigated. Low dehydroepiandrosterone levels have been associated with increased risk of ischemic stroke, with women in the lowest quartile having a relative risk (RR) of 1.41 (95% confidence interval [CI], 1.03–1.92) for ischemic stroke after adjustment for other risk factors.⁴ Dehydroepiandrosterone levels at presentation of acute stroke were also inversely associated with stroke severity in a hospital-based study of stroke in postmenopausal women.⁵ Another study of women undergoing coronary angiography provided evidence that lower dehydroepiandrosterone levels were associated with increased cardiovascular mortality, including death from stroke.⁶

Additional prospective studies of endogenous sex hormones and risk of stroke in women are needed, particularly with more sensitive measures of hormones, and in high-risk groups including black and Hispanic women.

Received October 11, 2017; final revision received November 28, 2017; accepted December 1, 2017.

From the Department of Neurology and Ophthalmology and Department of Pharmacology and Toxicology, Michigan State University, East Lansing (S.L.D.); Baltimore Veterans Administration Medical Center and Department of Neurology, University of Maryland School of Medicine (S.K.); Channing Division of Network Medicine and Division of Women's Health (K.M.R.), Department of Medicine, Brigham and Women's Hospital, Boston, MA (S.H.L.); Department of Nutrition, Harvard T. H. Chan School of Public Health, Boston, MA (S.H.L.); and University of Michigan Stroke Program, Ann Arbor (M.M.).

Correspondence to Kathryn M. Rexrode, MD, MPH, Division of Women's Health, Department of Medicine, Brigham and Women's Hospital, 75 Francis St, OBC3 Boston, MA 02115. E-mail krexrode@bwh.harvard.edu

(*Stroke*. 2018;49:00-00. DOI: 10.1161/STROKEAHA.117.018415.)

© 2018 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.117.018415

Age at Menarche

Earlier age at menarche has been associated with greater cardiovascular disease (CVD) morbidity and mortality in some,^{7–9} but not all studies,^{10–12} and data specific for stroke are limited. In the Million Women Study from the United Kingdom, a U-shaped relationship between age at menarche and cerebrovascular disease was observed.⁹ Women who experienced menarche at age ≤ 10 years were at higher risk of developing stroke in later life compared with those with age at menarche at 13 years (RR, 1.16 [95% CI, 1.09–1.23]); however, women who experienced menarche at age ≥ 17 years were also at higher risk of developing stroke compared with those with age at menarche at 13 years (RR, 1.13 [95% CI, 1.03–1.24]).⁹ Women with extremely early age at menarche may experience hormonal disturbances, such as higher exposure to estradiol, potentially mediated through childhood obesity.¹³ The timing of menarche is associated with type 2 diabetes mellitus risk,^{14,15} an association which may be influenced by childhood adiposity and endocrinopathies.^{13,16} Although the Million Women Study attempted to control for potential confounding factors, such as body mass index, through statistical adjustment,⁹ obesity may have been present before the onset of menarche, complicating inferences about cause and effect.

Age at Natural Menopause and Surgical Menopause

Women of reproductive age are at a lower risk of CVD compared with men of similar age and lifestyle, but women who experience early menopause have increased cardiovascular risk.¹⁷ In a recent meta-analysis, investigators reported that early age at natural menopause (menopause onset before 45 years) was associated with a slightly higher risk of total CVD mortality (RR, 1.12 [95% CI, 1.03–1.21]) than onset at age 45 years or later; however, this association was not observed for stroke mortality risk independently.¹⁸ Surgical menopause, bilateral oophorectomy with or without hysterectomy, has also been associated with higher risk of CVD.^{19,20} In the Nurses' Health Study, bilateral oophorectomy before age 50 years was associated with increased CVD mortality in women and especially in women who did not use hormone therapy.²¹ When a sensitivity analysis of stroke mortality was conducted, the CI of this association widened potentially because of low numbers although the risk estimate remained elevated (RR, 1.15 [95% CI, 0.85–1.56]).²¹ Therefore, further investigations are warranted to examine these potential associations for early menopause and stroke.

Specific mechanisms responsible for the association between the timing of age at menopause and CVD are unclear. However, CVD incidence rising sharply after menopause suggests protective benefits of ovarian hormones.²² Estrogen inhibits hepatic lipase,²³ thus decline in endogenous estrogens in the menopausal transition may adversely affect lipid levels and subsequently cardiovascular risk.²⁴ The menopausal transition is associated with declines in high-density lipoprotein cholesterol and increases in low-density lipoprotein cholesterol,²⁴ as well as changes in high-density lipoprotein composition, with higher number of small high-density lipoprotein particles, which confer less cardiovascular protection than large high-density lipoprotein particles.²⁵ Hence, decreased estrogen

concentrations over the menopause transition and effects on lipoprotein profiles may subsequently contribute to atherosclerosis. In a cross-sectional, population-based study, longer duration of reproductive lifespan and years from menarche to menopause, were associated with a lower 10-year CVD risk assessed by the Framingham Risk Score in postmenopausal women.²⁶ In the prospective cohort Nurses' Health Study, a shorter duration of reproductive lifespan was associated with a higher risk of stroke, as well as CVD, which was likely driven by earlier age at menopause (either naturally or surgically).²⁷

Exogenous Estrogens and Stroke Risk

Hormone-Containing Birth Control and Stroke

Hormonal contraceptives, including oral, transdermal, and vaginal formulations, are effective and are used worldwide by >100 million women (World Health Organization 2014). There are various formulations containing either combined estrogen and progestogen or progestogen alone, administered as pills, patches, and rings. Combined oral contraceptives (COC), comprised both estrogen and progestogen, are thrombogenic²⁸ and, historically, have been associated with increased risk of CVD.^{29,30} Oral estrogens have a dose–response association with risk, and doses have declined since their introduction in the 1960s. Most COCs now contain <50 μg and some contain as low as 15 μg of estrogen. Authors who evaluated the risk of second- and third-generation estrogen-containing oral contraceptives continued to find a 60% to 80% increased odds (95% CI, 1.2–2.8) of the combined end point of myocardial infarction or ischemic stroke among COC users compared with non-users.^{30–33} In a separate study, second-generation COCs were associated with an odds ratio=2.54 (95% CI, 1.96–3.28) and third-generation oral contraceptives with an odds ratio=2.03 (95% CI, 1.15–3.57) of stroke.³⁴ Progestogen-only hormonal contraceptives have not been associated with increased risk of ischemic stroke although data are limited.^{30,35} Nonoral methods of delivering combined hormonal contraceptives, including the vaginal ring and contraceptive patches, seem to have the same risk as oral contraceptives.³⁶

Risk of stroke with COC use rises in the presence of other cardiovascular risk factors (ie, smoking, age [>35 years], and history of migraine with aura). Migraine with aura is a common condition in younger women, and the risk of stroke in patients with migraine with aura is increased ≈ 2 -fold.³⁷ Women with migraine who also use COCs have a further increased risk of ischemic stroke (7.02 [95% CI, 1.51–32.68]) where women with migraine with aura, COC use, and who are active smokers have a dramatically elevated risk for stroke (RR, 10 [95% CI, 1.4–73.7]).³⁷ Guidelines from the International Headache Society Task Force on COC prescribing recommendations have been published previously.³⁸ Women who have migraine with aura should be advised to control all modifiable risk factors, including tobacco use and hypertension, and birth control methods other than COCs should be considered.³¹

Hormonal contraceptives are used by millions of women, and for most low-risk women, the risk of stroke associated with COC is lower than the risk of stroke during pregnancy. However, there is a clear association between hormone-containing birth control methods and ischemic stroke. This is

magnified by stroke risk factors. Although a COC pill containing 30 µg of estrogen is considered safe and effective hormonal contraception,³⁰ careful attention to stroke risk should be made before prescribing. Nonhormonal and progestogen-only methods of contraception should also be considered in high-risk patients. Further research to evaluate the risk of progestogen-only methods (depot injection, pills, implants, and intrauterine devices) is needed.

Postmenopausal Hormone Therapy and Stroke Risk

Prospective observational studies and randomized trials consistently demonstrate an increased risk of stroke, particularly ischemic stroke, with oral postmenopausal hormone therapy.

In prospective cohort studies, data suggest that postmenopausal users of oral estrogens with or without progestin have a 27% to 39% increased risk of stroke compared with nonusers.³⁹ In the Women's Health Initiative, women randomized to combined estrogen plus progestin had a hazard ratio of 1.31 (95% CI, 1.02–1.68) for total stroke and 1.44 (95% CI, 1.09–1.90) for ischemic stroke.⁴⁰ In the Women's Health Initiative trial of unopposed estrogen, women randomized to active therapy had a hazard ratio of 1.37 (95% CI, 1.09–1.73) for total stroke and 1.55 (95% CI, 1.19–2.01) for ischemic stroke.⁴¹

Although some data suggest an association between timing of hormone therapy and coronary heart disease, time since menopause is not associated with differences in stroke incidence in either observational studies³⁹ or clinical trials.⁴² The incidence of stroke is relatively low in younger women (age, 50–59 years), with ≈2 additional cases of stroke per 10000 women per year taking postmenopausal hormones.⁴² In addition, there is a dose-response relationship between dose of oral conjugated estrogen and stroke, with RRs of 0.93 for a dose of 0.3 mg, 1.54 at 0.625 mg, and 1.62 at 1.25 mg (*P* for trend, <0.001).³⁹

Similar results have been found in secondary cardiovascular prevention. Both the HERS (Heart and Estrogen/Progestin Replacement Study) in women with prior coronary heart disease⁴³ and the WEST (Women's Estrogen for Stroke Trial) in women with recent mild ischemic stroke or transient ischemic attack⁴⁴ found no significant effect of treatment with oral estrogen or combined oral estrogen and progestin on risk of stroke, with a trend toward harm.

Limited data are available for transdermal estrogens, which have been associated with lower risk of venous thromboembolism. In a population-based nested case-control study, current use of transdermal hormone therapy was not associated with an increased risk of stroke (hazard ratio, 0.95 [0.75–1.20]).⁴⁵ However, when the dose was examined, low-dose transdermal estrogen (≤50 µg/d estradiol) was not associated with risk while high-dose transdermal estrogen (>50 µg/d) was associated with increased stroke risk.⁴⁵

Transgender Medicine

Transgender individuals are people whose sex identity differs from their sex assigned at birth. The prevalence of self-identified transgender adults in the United States is estimated to be ≈0.5% of the population.⁴⁶ Some transgender people pursue hormonal therapy or sex-affirming surgery to assume secondary sex characteristics consistent with their sex identity. The

use of certain hormonal therapies has implications for the incidence of cerebrovascular disease in these individuals.

Transwomen are people with an assigned male sex and a female sex identity. Transwomen may undergo medical treatment with estrogens, antiandrogens, or a combination of both.⁴⁷ Those who have undergone orchiectomy may pursue only estrogen therapy.

Antiandrogen therapies do not seem to increase stroke risk in transwomen. Spironolactone is the antiandrogen most commonly prescribed to transwomen in the United States.⁴⁷ Spironolactone, a potassium-sparing diuretic, may lower blood pressure but does not increase thrombotic risk. Similarly, finasteride, a less commonly used antiandrogen, does not seem to increase thrombotic risk.

Direct data on the effect of exogenous estrogens in transwomen are scant. Much of our knowledge about the effects of exogenous estrogen is derived from studies of the increased risk of thrombotic complications, including stroke among postmenopausal women using postmenopausal hormone therapy.^{48,49} Prospective trials of thrombotic risk in transwomen receiving estrogen therapy are lacking.⁵⁰ A 1997 Dutch single-center retrospective descriptive study of 816 transwomen treated for a mean 9.5 patient-years with ethinyl estradiol and the antiandrogen cyproterone acetate found that 45 (5.5%) developed deep vein thrombosis or pulmonary embolism, 5 (0.6%) experienced a transient ischemic attack, and none experienced ischemic stroke.⁵¹ A 2013 Belgian single-center case-control study evaluated 214 transwomen who were maintained on estrogen therapy for a median of 6 years.⁵² Eleven of the 214 transwomen (5.1%) developed deep vein thrombosis or pulmonary embolism during hormonal treatment. Five of the 214 transwomen (2.3%) were diagnosed with transient ischemic attack or cerebrovascular disease during treatment, a higher prevalence than in the age-matched control men. In a 2011 Dutch retrospective single-clinic cohort mortality study of 966 transwomen on estrogen with or without antiandrogen therapy with a mean follow-up of 19.4 years, no difference was found in the incidence of fatal stroke in transwomen compared with the incidence in the general population.⁵³

As we await prospective studies of estrogen therapy in transwomen, we recommend that medical providers maintain a high index of suspicion for deep vein thrombosis/pulmonary embolism and cerebral venous thrombosis in transwomen receiving estrogen therapy. Cardiovascular risk should be evaluated, and transwomen who smoke should be encouraged to quit smoking and provided with appropriate pharmacological and psychosocial supports.

Transmen are individuals with an assigned female sex and male sex identity. Transmen may undergo treatment with testosterone to promote development of male secondary sex characteristics. Testosterone is available via transdermal and intramuscular routes. Unlike estrogen, testosterone does not seem to be associated with an increased risk of thromboembolic complications. The majority of existing studies of transmen do not suggest an increased risk of cardiovascular morbidity with exogenous testosterone therapy.^{51–54}

In general, providers should be aware that transgender people may be less likely to seek medical attention than their cisgender peers because of previous negative interactions with the

Table. Summary of Female-Specific Stroke Risk Factors

Exposure	Risk Association	Further Research Needed
Endogenous hormones		
Early age at menarche (<10 y)	↑	
Early age at menopause/BSO (<45 y)	↑	
Reproductive lifespan	?	Yes
Low DHEAS	↑	
Estradiol	?	Yes
Testosterone	→	
Exogenous hormones		
PMH: oral estrogens	↑	
PMH: transdermal estrogens	?	Yes
Combined oral contraceptives	↑	
Progestogen-only contraceptives	→	Yes
Transgender exogenous estrogens	↑	Yes
Transgender exogenous testosterone	→	Yes
Pregnancy-related exposures		
Pregnancy/periparturition	↑	
Gestational diabetes mellitus	↑	
Hypertension in pregnancy/preeclampsia	↑	Yes

BSO indicates bilateral salpingo-oophorectomy; DHEAS, dehydroepiandrosterone; and PMH, postmenopausal hormones.

medical community, poor psychosocial wellbeing, and fear of stigma.^{55,56} This has direct implications for delay of appropriate stroke care. Health systems and medical providers can improve the medical care of transgender people by asking about sex identity and preferred pronouns on intake forms⁵⁷; promoting an explicitly supportive and inclusive clinic or unit culture for transgender patients and their families; and pursuing research into the unique health needs of transmen and transwomen.

Issues of Pregnancy, Parity, and Stroke

Pregnancy and the peripartum are associated with increased risk of stroke. The peripartum period from 2 days before to 1 day after delivery and, to a lesser extent, up to 6 weeks postpartum is associated with an increased risk of ischemic stroke and intracerebral hemorrhage.^{58–60} In a large population-based study of women in England (age, 15–49 years), authors found that the baseline incidence of stroke was 25.0/100 000 person-years in women when they were not pregnant. The incidence rate dropped during early pregnancy but was 9-fold higher in the peripartum period (161.1/100 000 person-years) and 3-fold higher in the early postpartum period (47.1/100 000 person-years; 95% CI, 31.3–70.9).⁶¹ For subarachnoid hemorrhage, only the peripartum period confers increased risk^{60,61}; nonaneurysmal subarachnoid hemorrhage is likely a major contributor to this risk. The risk of any thrombotic event that includes ischemic stroke remains increased to a lesser extent until 12 weeks postpartum,⁶² but it is not established that the risk of stroke remains increased

beyond 6 weeks postpartum.⁶¹ Eclampsia and preeclampsia are the strongest risk factors for both ischemic stroke and intracerebral hemorrhage accounting for 24% to 48% of all pregnancy-associated strokes^{58,59}; this risk is potentiated by preexisting genitourinary tract infection, chronic hypertension, prothrombotic states, and coagulopathies.⁶³

Complications of pregnancy, specifically pregnancy-induced hypertension, gestational diabetes mellitus, and preeclampsia are also associated with long-term risk of stroke.⁶⁴ There is evidence that women with pregnancy outcomes of preterm birth and small for gestational age infants have higher rates of cerebrovascular events even after adjusting for other pregnancy complications.⁶⁵

For women with prior stroke, the risk of recurrent stroke is increased in the peripartum and postpartum periods. Limited data from case series^{66–68} suggest an absolute risk of recurrent arterial ischemic stroke associated with pregnancy of 0.7%, similar to the <1% yearly risk of recurrent stroke among young adults who have no vascular risk factors⁶⁹; however, the 95% CI is wide, 0.04% to 4.4%, indicating the need for further study. In addition, the absolute risk depends on clinical circumstances, with the presence of vascular risk factors or a definite cause of stroke, including thrombophilic disorders, conferring an increased risk. Similarly, there is a paucity of information on the excess risk of pregnancy complications to mother and child among women with prior ischemic stroke.⁷⁰

The role of pregnancy on risk of intracranial hemorrhage from preexisting arteriovenous malformations is uncertain. The 2 largest case-crossover studies of arteriovenous malformations, single-center studies from China⁷¹ and Baltimore,⁷² yielded conflicting results. The preponderance of available evidence suggests that pregnancy and delivery does not increase the risk of aneurysm rupture.^{73,74}

Conclusions

It is important to be aware of stroke risk factors specific to women. The Table summarizes the associations for stroke risk factors unique to women. Specific considerations that include endogenous hormone levels, exogenous hormone therapy, pregnancy and the peripartum period, and pregnancy-related complications change the risk of stroke for women as well as the optimal stroke prevention strategies. Special attention and further research are also needed in the area of transgender individuals. Further research to determine whether risk prediction models should include risk factors specific to women, including hormonal and reproductive exposures, is needed.

Disclosures

Dr Rexrode is currently supported by National Institutes of Health grant HL088521; Dr Demel receives compensation as a member of the Speaker's Bureau for Genentech. The other authors report no conflicts.

References

- Holmegard HN, Nordestgaard BG, Jensen GB, Tybjaerg-Hansen A, Benn M. Sex hormones and ischemic stroke: a prospective cohort study and meta-analyses. *J Clin Endocrinol Metab*. 2016;101:69–78. doi: 10.1210/jc.2015-2687.
- Scarabin-Carré V, Canonico M, Brailly-Tabard S, Trabado S, Ducimetière P, Giroud M, et al. High level of plasma estradiol as a new predictor

- of ischemic arterial disease in older postmenopausal women: the three-city cohort study. *J Am Heart Assoc.* 2012;1:e001388. doi: 10.1161/JAHA.112.001388.
3. Lee JS, Yaffe K, Lui LY, Cauley J, Taylor B, Browner W, et al; Study of Osteoporotic Fractures Group. Prospective study of endogenous circulating estradiol and risk of stroke in older women. *Arch Neurol.* 2010;67:195–201. doi: 10.1001/archneurol.2009.322.
 4. Jiménez MC, Sun Q, Schürks M, Chiuvè S, Hu FB, Manson JE, et al. Low dehydroepiandrosterone sulfate is associated with increased risk of ischemic stroke among women. *Stroke.* 2013;44:1784–1789. doi: 10.1161/STROKEAHA.111.000485.
 5. Pappa T, Vemmos K, Saltiki K, Mantzou E, Stamatelopoulos K, Alevizaki M. Severity and outcome of acute stroke in women: relation to adrenal sex steroid levels. *Metabolism.* 2012;61:84–91. doi: 10.1016/j.metabol.2011.06.003.
 6. Shufelt C, Bretsky P, Almeida CM, Johnson BD, Shaw LJ, Azziz R, et al. DHEA-S levels and cardiovascular disease mortality in postmenopausal women: results from the National Institutes of Health–National Heart, Lung, and Blood Institute (NHLBI)-sponsored Women’s Ischemia Syndrome Evaluation (WISE). *J Clin Endocrinol Metab.* 2010;95:4985–4992. doi: 10.1210/jc.2010-0143.
 7. Cooper GS, Ephross SA, Weinberg CR, Baird DD, Whelan EA, Sandler DP. Menstrual and reproductive risk factors for ischemic heart disease. *Epidemiology.* 1999;10:255–259.
 8. Jacobsen BK, Oda K, Knutsen SF, Fraser GE. Age at menarche, total mortality and mortality from ischaemic heart disease and stroke: the Adventist Health Study, 1976–88. *Int J Epidemiol.* 2009;38:245–252. doi: 10.1093/ije/dyn251.
 9. Canoy D, Beral V, Balkwill A, Wright FL, Kroll ME, Reeves GK, et al; Million Women Study Collaborators*. Age at menarche and risks of coronary heart and other vascular diseases in a large UK cohort. *Circulation.* 2015;131:237–244. doi: 10.1161/CIRCULATIONAHA.114.010070.
 10. Colditz GA, Willett WC, Stampfer MJ, Rosner B, Speizer FE, Hennekens CH. A prospective study of age at menarche, parity, age at first birth, and coronary heart disease in women. *Am J Epidemiol.* 1987;126:861–870.
 11. Charalampopoulos D, McLoughlin A, Elks CE, Ong KK. Age at menarche and risks of all-cause and cardiovascular death: a systematic review and meta-analysis. *Am J Epidemiol.* 2014;180:29–40. doi: 10.1093/aje/kwu113.
 12. Cui R, Iso H, Toyoshima H, Date C, Yamamoto A, Kikuchi S, et al; JACC Study Group. Relationships of age at menarche and menopause, and reproductive year with mortality from cardiovascular disease in Japanese postmenopausal women: the JACC study. *J Epidemiol.* 2006;16:177–184.
 13. Zhai L, Liu J, Zhao J, Liu J, Bai Y, Jia L, et al. Association of obesity with onset of puberty and sex hormones in Chinese girls: a 4-year longitudinal study. *PLoS One.* 2015;10:e0134656. doi: 10.1371/journal.pone.0134656.
 14. He C, Zhang C, Hunter DJ, Hankinson SE, Buck Louis GM, Hediger ML, et al. Age at menarche and risk of type 2 diabetes: results from 2 large prospective cohort studies. *Am J Epidemiol.* 2010;171:334–344. doi: 10.1093/aje/kwp372.
 15. Elks CE, Ong KK, Scott RA, van der Schouw YT, Brand JS, Wark PA, et al; InterAct Consortium. Age at menarche and type 2 diabetes risk: the EPIC-InterAct study. *Diabetes Care.* 2013;36:3526–3534. doi: 10.2337/dc13-0446.
 16. Pinkney J, Streeter A, Hosking J, Mostazir M, Mohammad M, Jeffery A, et al. Adiposity, chronic inflammation, and the prepubertal decline of sex hormone binding globulin in children: evidence for associations with the timing of puberty (Earlybird 58). *J Clin Endocrinol Metab.* 2014;99:3224–3232. doi: 10.1210/jc.2013-3902.
 17. Atsma F, Bartelink ML, Grobbee DE, van der Schouw YT. Postmenopausal status and early menopause as independent risk factors for cardiovascular disease: a meta-analysis. *Menopause.* 2006;13:265–279. doi: 10.1097/01.gme.0000218683.97338.ea.
 18. Muka T, Oliver-Williams C, Kunutsor S, Laven JS, Fauser BC, Chowdhury R, et al. Association of age at onset of menopause and time since onset of menopause with cardiovascular outcomes, intermediate vascular traits, and all-cause mortality: a systematic review and meta-analysis. *JAMA Cardiol.* 2016;1:767–776. doi: 10.1001/jamacardio.2016.2415.
 19. Parker WH, Broder MS, Chang E, Feskanich D, Farquhar C, Liu Z, et al. Ovarian conservation at the time of hysterectomy and long-term health outcomes in the nurses’ health study. *Obstet Gynecol.* 2009;113:1027–1037. doi: 10.1097/AOG.0b013e3181a11c64.
 20. Howard BV, Kuller L, Langer R, Manson JE, Allen C, Assaf A, et al; Women’s Health Initiative. Risk of cardiovascular disease by hysterectomy status, with and without oophorectomy: the Women’s Health Initiative Observational Study. *Circulation.* 2005;111:1462–1470. doi: 10.1161/01.CIR.0000159344.21672.FD.
 21. Parker WH, Feskanich D, Broder MS, Chang E, Shoupe D, Farquhar CM, et al. Long-term mortality associated with oophorectomy compared with ovarian conservation in the nurses’ health study. *Obstet Gynecol.* 2013;121:709–716. doi: 10.1097/AOG.0b013e3182864350.
 22. Vitale C, Fini M, Speziale G, Chierchia S. Gender differences in the cardiovascular effects of sex hormones. *Fundam Clin Pharmacol.* 2010;24:675–685. doi: 10.1111/j.1472-8206.2010.00817.x.
 23. Berg GA, Siseles N, González AI, Ortiz OC, Tempone A, Wikinski RW. Higher values of hepatic lipase activity in postmenopause: relationship with atherogenic intermediate density and low density lipoproteins. *Menopause.* 2001;8:51–57.
 24. Matthews KA, Meilahn E, Kuller LH, Kelsey SF, Caggiula AW, Wing RR. Menopause and risk factors for coronary heart disease. *N Engl J Med.* 1989;321:641–646. doi: 10.1056/NEJM198909073211004.
 25. Woodard GA, Brooks MM, Barinas-Mitchell E, Mackey RH, Matthews KA, Sutton-Tyrrell K. Lipids, menopause, and early atherosclerosis in Study of Women’s Health Across the Nation Heart women. *Menopause.* 2011;18:376–384. doi: 10.1097/gme.0b013e3181f6480e.
 26. Kim SH, Sim MY, Park SB. Association between duration of reproductive lifespan and Framingham risk score in postmenopausal women. *Maturitas.* 2015;82:431–435. doi: 10.1016/j.maturitas.2015.07.011.
 27. Ley SH, Li Y, Tobias DK, Manson JE, Rosner B, Hu FB, et al. Duration of reproductive life span, age at menarche, and age at menopause are associated with risk of cardiovascular disease in women. *J Am Heart Assoc.* 2017;6:e006713:1-11. doi: 10.1161/JAHA.117.006713.
 28. Tchaikovski SN, Rosing J. Mechanisms of estrogen-induced venous thromboembolism. *Thromb Res.* 2010;126:5–11. doi: 10.1016/j.thromres.2010.01.045.
 29. Zakharova MY, Meyer RM, Brandy KR, Datta YH, Joseph MS, Schreiner PJ, et al. Risk factors for heart attack, stroke, and venous thrombosis associated with hormonal contraceptive use. *Clin Appl Thromb Hemost.* 2011;17:323–331. doi: 10.1177/1076029610368670.
 30. Roach RE, Helmerhorst FM, Lijfering WM, Stijnen T, Algra A, Dekkers OM. Combined oral contraceptives: the risk of myocardial infarction and ischemic stroke. *Cochrane Database Syst Rev.* 2015;CD011054. doi: 10.1002/14651858.CD011054.pub2.
 31. Plu-Bureau G, Hugon-Rodin J, Maitrot-Mantelet L, Canonico M. Hormonal contraceptives and arterial disease: an epidemiological update. *Best Pract Res Clin Endocrinol Metab.* 2013;27:35–45. doi: 10.1016/j.beem.2012.11.003.
 32. Chan WS, Ray J, Wai EK, Ginsburg S, Ginsburg S, Hannah ME, et al. Risk of stroke in women exposed to low-dose oral contraceptives: a critical evaluation of the evidence. *Arch Intern Med.* 2004;164:741–747. doi: 10.1001/archinte.164.7.741.
 33. Gillum LA, Mamidipudi SK, Johnston SC. Ischemic stroke risk with oral contraceptives: a meta-analysis. *JAMA.* 2000;284:72–78.
 34. Baillargeon JP, McClish DK, Essah PA, Nestler JE. Association between the current use of low-dose oral contraceptives and cardiovascular arterial disease: a meta-analysis. *J Clin Endocrinol Metab.* 2005;90:3863–3870. doi: 10.1210/jc.2004-1958.
 35. Chakhtoura Z, Canonico M, Gompel A, Thalabard JC, Scarabin PY, Plu-Bureau G. Progestogen-only contraceptives and the risk of stroke: a meta-analysis. *Stroke.* 2009;40:1059–1062. doi: 10.1161/STROKEAHA.108.538405.
 36. Lidegaard Ø, Løkkegaard E, Jensen A, Skovlund CW, Keiding N. Thrombotic stroke and myocardial infarction with hormonal contraception. *N Engl J Med.* 2012;366:2257–2266. doi: 10.1056/NEJMoa1111840.
 37. Schürks M, Rist PM, Bigal ME, Buring JE, Lipton RB, Kurth T. Migraine and cardiovascular disease: systematic review and meta-analysis. *BMJ.* 2009;339:b3914.
 38. Bousser MG, Conard J, Kittner S, de Lignières B, MacGregor EA, Massiou H, et al. Recommendations on the risk of ischaemic stroke associated with use of combined oral contraceptives and hormone replacement therapy in women with migraine. The International Headache Society Task Force on Combined Oral Contraceptives & Hormone Replacement Therapy. *Cephalalgia.* 2000;20:155–156. doi: 10.1046/j.1468-2982.2000.00035.x.
 39. Grodstein F, Manson JE, Stampfer MJ, Rexrode K. Postmenopausal hormone therapy and stroke: role of time since menopause and age at

- initiation of hormone therapy. *Arch Intern Med*. 2008;168:861–866. doi: 10.1001/archinte.168.8.861.
40. Wassertheil-Smolter S, Hendrix SL, Limacher M, Heiss G, Kooperberg C, Baird A, et al; WHI Investigators. Effect of estrogen plus progestin on stroke in postmenopausal women: the Women's Health Initiative: a randomized trial. *JAMA*. 2003;289:2673–2684. doi: 10.1001/jama.289.20.2673.
 41. Hendrix SL, Wassertheil-Smolter S, Johnson KC, Howard BV, Kooperberg C, Rossouw JE, et al; WHI Investigators. Effects of conjugated equine estrogen on stroke in the Women's Health Initiative. *Circulation*. 2006;113:2425–2434. doi: 10.1161/CIRCULATIONAHA.105.594077.
 42. Rossouw JE, Prentice RL, Manson JE, Wu L, Barad D, Barnabei VM, et al. Postmenopausal hormone therapy and risk of cardiovascular disease by age and years since menopause. *JAMA*. 2007;297:1465–1477. doi: 10.1001/jama.297.13.1465.
 43. Simon JA, Hsia J, Cauley JA, Richards C, Harris F, Fong J, et al. Postmenopausal hormone therapy and risk of stroke: the Heart and Estrogen/progestin Replacement Study (HERS). *Circulation*. 2001;103:638–642.
 44. Viscoli CM, Brass LM, Kernan WN, Sarrel PM, Suissa S, Horwitz RI. A clinical trial of estrogen-replacement therapy after ischemic stroke. *N Engl J Med*. 2001;345:1243–1249. doi: 10.1056/NEJMoa010534.
 45. Renoux C, Dell'aniello S, Garbe E, Suissa S. Transdermal and oral hormone replacement therapy and the risk of stroke: a nested case-control study. *BMJ*. 2010;340:c2519.
 46. Crissman HP, Berger MB, Graham LF, Dalton VK. Transgender demographics: a household probability sample of US adults, 2014. *Am J Public Health*. 2017;107:213–215. doi: 10.2105/AJPH.2016.303571.
 47. Chipkin SR, Kim F. Ten most important things to know about caring for transgender patients. *Am J Med*. 2017;130:1238–1245. doi: 10.1016/j.amjmed.2017.06.019.
 48. Canonico M, Plu-Bureau G, Lowe GD, Scarabin PY. Hormone replacement therapy and risk of venous thromboembolism in postmenopausal women: systematic review and meta-analysis. *BMJ*. 2008;336:1227–1231.
 49. Boardman HM, Hartley L, Eisinga A, Main C, Roque i Figuls M, Bonfill Cosp X, et al. Hormone therapy for preventing cardiovascular disease in post-menopausal women. *Cochrane Database Syst Rev*. 2015;Cd002229. doi: 10.1002/14651858.CD002229.pub4.
 50. Shatzel JJ, Connelly KJ, DeLoughery TG. Thrombotic issues in transgender medicine: a review. *Am J Hematol*. 2017;92:204–208. doi: 10.1002/ajh.24593.
 51. van Kesteren PJ, Asscheman H, Megens JA, Gooren LJ. Mortality and morbidity in transsexual subjects treated with cross-sex hormones. *Clin Endocrinol (Oxf)*. 1997;47:337–342.
 52. Wierckx K, Elaut E, Declercq E, Heylens G, De Cuypere G, Taes Y, et al. Prevalence of cardiovascular disease and cancer during cross-sex hormone therapy in a large cohort of trans persons: a case-control study. *Eur J Endocrinol*. 2013;169:471–478. doi: 10.1530/EJE-13-0493.
 53. Asscheman H, Giltay EJ, Megens JA, de Ronde WP, van Trotsenburg MA, Gooren LJ. A long-term follow-up study of mortality in transsexuals receiving treatment with cross-sex hormones. *Eur J Endocrinol*. 2011;164:635–642. doi: 10.1530/EJE-10-1038.
 54. Gooren LJ, Wierckx K, Giltay EJ. Cardiovascular disease in transsexual persons treated with cross-sex hormones: reversal of the traditional sex difference in cardiovascular disease pattern. *Eur J Endocrinol*. 2014;170:809–819. doi: 10.1530/EJE-14-0011.
 55. Winter S, Diamond M, Green J, Karasic D, Reed T, Whittle S, et al. Transgender people: health at the margins of society. *Lancet*. 2016;388:390–400. doi: 10.1016/S0140-6736(16)00683-8.
 56. Samuels EA, Tape C, Garber N, Bowman S, Choo EK. "Sometimes you feel like the freak show": a qualitative assessment of emergency care experiences among transgender and gender-nonconforming patients [published online ahead of print July 13, 2017]. *Ann Emerg Med*. 2017. doi: 10.1016/j.annemergmed.2017.05.002. <http://www.sciencedirect.com/science/article/pii/S0196064417305838?via%3Dihub>.
 57. Schuster MA, Reisner SL, Onorato SE. Beyond bathrooms—meeting the health needs of transgender people. *N Engl J Med*. 2016;375:101–103. doi: 10.1056/NEJMp1605912.
 58. Sharshar T, Lamy C, Mas JL. Incidence and causes of strokes associated with pregnancy and puerperium. A study in public hospitals of Ile de France. Stroke in Pregnancy Study Group. *Stroke*. 1995;26:930–936.
 59. Kittner SJ, Stern BJ, Feeser BR, Hebel R, Nagey DA, Buchholz DW, et al. Pregnancy and the risk of stroke. *N Engl J Med*. 1996;335:768–774. doi: 10.1056/NEJM199609123351102.
 60. Salonen Ros H, Lichtenstein P, Bellocco R, Petersson G, Cnattingius S. Increased risks of circulatory diseases in late pregnancy and puerperium. *Epidemiology*. 2001;12:456–460.
 61. Ban L, Sprigg N, Abdul Sultan A, Nelson-Piercy C, Bath PM, Ludvigsson JF, et al. Incidence of first stroke in pregnant and nonpregnant women of childbearing age: a population-based cohort study from England. *J Am Heart Assoc*. 2017;6:e004601:1–9. doi: 10.1011/JAHA.116.004601.
 62. Kamel H, Navi BB, Sriram N, Hovsepian DA, Devereux RB, Elkind MS. Risk of a thrombotic event after the 6-week postpartum period. *N Engl J Med*. 2014;370:1307–1315. doi: 10.1056/NEJMoa1311485.
 63. Miller EC, Gatollari HJ, Too G, Boehme AK, Leffert L, Marshall RS, et al. Risk factors for pregnancy-associated stroke in women with preeclampsia. *Stroke*. 2017;48:1752–1759. doi: 10.1161/STROKEAHA.117.017374.
 64. Bushnell C, McCullough LD, Awad IA, Chireau MV, Fedder WN, Furie KL, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Epidemiology and Prevention; Council for High Blood Pressure Research. Guidelines for the prevention of stroke in women: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:1545–1588. doi: 10.1161/01.str.0000442009.06663.48.
 65. Bonamy AK, Parikh NI, Cnattingius S, Ludvigsson JF, Ingelsson E. Birth characteristics and subsequent risks of maternal cardiovascular disease: effects of gestational age and fetal growth. *Circulation*. 2011;124:2839–2846. doi: 10.1161/CIRCULATIONAHA.111.034884.
 66. Lamy C, Hamon JB, Coste J, Mas JL. Ischemic stroke in young women: risk of recurrence during subsequent pregnancies. French Study Group on Stroke in Pregnancy. *Neurology*. 2000;55:269–274.
 67. Coppage KH, Hinton AC, Moldenbauer J, Kovilam O, Barton JR, Sibai BM. Maternal and perinatal outcome in women with a history of stroke. *Am J Obstet Gynecol*. 2004;190:1331–1334. doi: 10.1016/j.ajog.2003.11.002.
 68. Crovetto F, Ossola MW, Spadaccini G, Duiella SF, Somigliana E, Fedele L. Ischemic stroke recurrence during pregnancy: a case series and a review of the literature. *Arch Gynecol Obstet*. 2012;286:599–604. doi: 10.1007/s00404-012-2352-0.
 69. Putaala J, Haapaniemi E, Metso AJ, Metso TM, Arto V, Kaste M, et al. Recurrent ischemic events in young adults after first-ever ischemic stroke. *Ann Neurol*. 2010;68:661–671. doi: 10.1002/ana.22091.
 70. Aarnio K GM, Gissler M, Grittner U, Siegerink B, Kaste M, Tatlisumak T, et al. Outcome of pregnancies and deliveries before and after ischaemic stroke. *European Stroke J*. 2017;26:346–355.
 71. Liu XJ, Wang S, Zhao YL, Teo M, Guo P, Zhang D, et al. Risk of cerebral arteriovenous malformation rupture during pregnancy and puerperium. *Neurology*. 2014;82:1798–1803. doi: 10.1212/WNL.0000000000000436.
 72. Porras JL, Yang W, Philadelphia E, Law J, Garzon-Muvdi T, Caplan JM, et al. Hemorrhage risk of brain arteriovenous malformations during pregnancy and puerperium in a North American Cohort. *Stroke*. 2017;48:1507–1513. doi: 10.1161/STROKEAHA.117.016828.
 73. Tiel Groenestege AT, Rinkel GJ, van der Bom JG, Algra A, Klijn CJ. The risk of aneurysmal subarachnoid hemorrhage during pregnancy, delivery, and the puerperium in the Utrecht population: case-crossover study and standardized incidence ratio estimation. *Stroke*. 2009;40:1148–1151. doi: 10.1161/STROKEAHA.108.539700.
 74. Kim YW, Neal D, Hoh BL. Cerebral aneurysms in pregnancy and delivery: pregnancy and delivery do not increase the risk of aneurysm rupture. *Neurosurgery*. 2013;72:143–149; discussion 150. doi: 10.1227/NEU.0b013e3182796af9.

KEY WORDS: contraception ■ estrogens ■ postmenopause ■ pregnancy ■ stroke

Stroke

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Stroke Risk Factors Unique to Women

Stacie L. Demel, Steven Kittner, Sylvia H. Ley, Mollie McDermott and Kathryn M. Rexrode

Stroke. published online February 8, 2018;

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2018 American Heart Association, Inc. All rights reserved.

Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/early/2018/02/07/STROKEAHA.117.018415.citation>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Stroke* is online at:
<http://stroke.ahajournals.org/subscriptions/>